



Influence of Long Term Fertilization and Manures on Soil Physical and Chemical Properties of Udorthentic Chromusterts

V. Venkatesh ^{a*#}, N. Chandra Sekaran ^{a†}, V. Sanjivkumar ^{b‡}, S. Meena ^{a†}
and K. T. Parthiban ^{c†}

^a Department of Soil Science and Agricultural Chemistry, AC&RI, Coimbatore-641003, India.

^b ARS, Kovilpatti- 628501, India.

^c Department of Agro-forestry, FC&RI, Mettupalayam-641301, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2021/v33i2430772

Editor(s):

(1) Dr. Hon H. Ho, State University of New York, USA.

Reviewers:

(1) Md Mozammel Haque, Bangladesh Rice Research Institute, Bangladesh.

(2) Bogdan-Vasile Cioruța, Technical University of Cluj-Napoca, Romania.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/79880>

Original Research Article

Received 06 October 2021
Accepted 12 December 2021
Published 14 December 2021

ABSTRACT

To study the effect of different nutrient management practices on different soil physical and chemical properties in the permanent manurial experiment field of Tamil Nadu Agricultural University, which was established during 1982 at Agriculture Research Station, Kovilpatti. Soil physical and chemical properties are mainly affected by the continuous application of fertilizers or manures from years together. To study the above mentioned properties of soil the soil samples were collected from the permanent manurial experiment of kovilpatti where the Randomized Block Design (RBD) was followed with nine different treatments viz., T₁- Control; T₂- 100 % RDF (40:20:40 NPK kg ha⁻¹); T₃- 50% RDF (20:10:20 NPK kg ha⁻¹); T₄- 50% N (Crop residues); T₅- 50 % N (FYM); T₆- 50 % Inorganic N+ 50% organic N (crop residues) + P (50%) + K (50%) ; T₇- 50 % Inorganic N+ 50% organic N (FYM) + P (50%) + K (50%); T₈- 100 % RDF + 25 kg ZnSO₄ ha⁻¹; T₉- FYM - 12.5 t ha⁻¹. The effect of these treatments along with the depth (0-15 cm; 15-30 cm and 30-

Resesach Scholar;

† Professor;

‡ Assistan Professor;

*Corresponding author: E-mail: vunnamvenkatesh010694@gmail.com;

45 cm) was compared. The treatment receiving organics viz., T₉- FYM - 12.5 t ha⁻¹ was observed to be the best in all the physical and chemical properties which was then followed by INM viz., T₇- 50 % Inorganic N+ 50% organic N (FYM) + P (50%) + K (50%) and T₆- 50 % Inorganic N+ 50% organic N (crop residues) + P (50%) + K (50%).

Keywords: Permanent manurial experiment; physical properties; chemical properties.

1. INTRODUCTION

Cotton is one of the major commercial crops that is been cultivated all over the world. The textile industry's primary source of raw materials is cotton. In India, the total cultivated area is 129.57 lakh ha with a production value of 371 lakh bales and productivity of 486.76 kg ha⁻¹ (As estimated by the Committee on Cotton Production and Consumption (COCP) in its meeting held on 25.01.2021). In Tamil Nadu, it is cultivated in an area of 1.31 lakh ha with a production of 6.00 lakh bales and productivity of 778.63 kg ha⁻¹. (Source: Cotton Advisory Board (CAB) P-Provisional as estimated by CAB on 18.6.2019).

Cotton's growth and yield are influenced by the interaction of the environment with the genetic makeup of the variety or hybrid, as well as a variety of inputs such as water, fertilizer, pesticides, and other factors. Fertilizers are one of the most influential factors in plant growth and development. Micronutrients and growth promoters are applied as foliar feeding, while primary nutrients such as N, P, and K are usually delivered to the crop through the soil.

The use of chemical fertilizers to enhance crop productivity has often negatively affected the complex system of the biogeochemical cycles [1,2] due to their continued long-term use. The overall strategy for increasing crop yields on a sustainable basis could be the conjunctive use of organic and inorganic sources of nutrients, along with other complementary measures. Organics are known to have favorable effects on soil structure, texture, and tilth and facilitate quick and greater availability of plant nutrients. Organics thus provide a better environment for root growth and proliferation, thereby creating a more absorptive surfaces for uptake of nutrients [3]. Some researchers have reported that integrated use of organic and inorganic sources of nutrients along with biofertilizers resulted in greater productivity, fertility, and net returns in soybean [4,5].

Wu et al. [6] reported that microbial inoculants increased the growth and nutritional assimilation

of total nitrogen (N), phosphorus (P), and potassium (K)] of maize and improved soil properties. Singh [7] found an increase in organic carbon and microbial biomass carbon in the treatments receiving the application of organic manures (particularly FYM), green manure, and bio-fertilizers in conjunction with inorganic fertilizer. More and Hangarge [8] noticed that grain and fodder yields of sorghum were greater in treatments receiving nutrients only through organics such as FYM, crop residues, and inoculation with azotobacter compared to nutrient supply only through chemical fertilizer. Studying the long-term effect of INM practices in terms of soil quality is of presumable importance in rainfed agriculture. With the above mention reports it is evident that INM showed an effect on different soil properties. Hence, the present study was mainly focused on the influence of continuous fertilization and manuring for a long term on soil physical and chemical properties.

2. MATERIALS AND METHODS

2.1 Site Description and Soil Analysis

In the year 1982, the Agricultural Research Station, Kovilpatti, located in the southern zone of Tamil Nadu (9.20' N altitude, 77.87' E longitude, 90m above MSL), began a permanent manurial experiment (PME) in the Cotton (KC 3) cropping system viz., cotton-fallow-fallow. With an annual mean rainfall of 743 mm and evapotranspiration of 812mm, the area is classified as a hot semi-arid region. The experimental site's soil falls under Kalathur soil series with fine montmorillonite, isohyperthermic, Udorthentic Chromusterts with heavy clay texture (Table 1). Soil samples were collected from each treatment followed for the past 30 years (Table 2) at three different depths of 0-15, 15-30, and 30-45cm by following quadrant method for the assessment of soil quality index in the Permanent manurial experiment. The samples collected were shade dried, ground with a wooden hammer, and passed through a 2 mm sieve. Finally, 1 kg of representative sample was preserved in a labeled cloth bag for laboratory analysis.

Table 1. Initial soil properties of experimental plot of Permanent manurial experiment in 1992

Properties	Value
EC	0.49 ds m ⁻¹
pH	8.2
Organic carbon	1.5 g kg ⁻¹
N	80 kg ha ⁻¹
P	10 kg ha ⁻¹
K	586 kg ha ⁻¹
Available Zn	1.2 Kg ha ⁻¹
Bulk density	1.23 to 1.30 Mg m ⁻³
Particle density	1.60 to 1.69 Mg m ⁻³
Infiltration rate	0.7 to 0.9 cm hr ⁻¹
Percentage of pore space	48.3 to 48.9
Permanent wilting point	14%
Field capacity	35 %
Coarse sand	10.90 to 11.50 %
Fine sand	9.45 to 14.10 %
Silt	15.6 to 19.95 %
Clay	48.05 to 53.00 %
Texture	Clay

Table 2. Permanent Manurial Experiment – Treatment details

T ₁	Control
T ₂	100 % RDF (40:20:40 NPK kg ha ⁻¹)
T ₃	50% RDF (20:10:20 NPK kg ha ⁻¹)
T ₄	50% N (Crop residues)
T ₅	50 % N (FYM)
T ₆	50 % Inorganic N+ 50% organic N (crop residues) + P (50%) + K (50%)
T ₇	50 % Inorganic N+ 50% organic N (FYM) + P (50%) + K (50%)
T ₈	100 % RDF + 25 kg ZnSO ₄ ha ⁻¹
T ₉	FYM - 12.5 t ha ⁻¹

*SSP- Single super Phosphate; FYM- Farm Yard Manure

2.2 Soil Analysis

2.2.1 Soil physical properties

The samples were analyzed for their physical properties viz., texture, Bulk density(BD), Particle density(PD), porosity, infiltration rate (IR), wet aggregate stability (WAS), Dry aggregate stability (DAS), and hydraulic conductivity as mentioned in Table 3.

2.2.2 Soil chemical properties

The chemical properties viz., Soil reaction (pH), Electrical Conductivity (EC), Cation exchange

capacity (CEC), Organic Carbon (OC), Nitrogen (N), Phosphorus (p), potassium (K) of the soil sample were analyzed by following the standard procedure as mentioned in Table 4.

2.3 Statistical Analysis

Randomized Block Design (RBD) was used for the experiment and the difference among the treatments was compared by the Least Significant Difference (LSD) test at a significance level of $p < 0.05$ using OPISTAT according to Gomez and Gomez [16].

Table 3. Details of analytical methods for soil physical properties analysis

S.No	Parameter	Methods adopted	Reference
1.	Bulk density, and pore space	Core sampler method	Gupta & Dakshinamoorthi [9]
2.	Hydraulic Conductivity	Constant hydraulic head method	Richards [10]

Table 4. Details of analytical methods for soil Chemical properties analysis

S.No	Parameter	Methods adopted	Reference
1.	Soil reaction (pH)	1:2.5 soil water extract	Jackson [11]
2.	Electrical conductivity	1:2.5 soil water extract	Jackson [11]
3.	Organic carbon	Chromic acid wet digestion	Walkley and Black [12]
4.	Available Nitrogen (KMnO ₄ – N)	Alkaline permanganate method	Subbiah and Asijia [13]
5.	Available phosphorus (Olsen – P)	Extraction with 0.5M NaHCO ₃	Olsen et al. [14]
6.	Available potassium (NH ₄ OA _C – K)	Extraction with neutral normal ammonium acetate and Flame photometry	Stanford and English [15]

3. RESULTS AND DISCUSSION

The soil samples collected from PME of Kovilpatti were analyzed for different physical (Table 5,6) and chemical (Table 7,8) properties, and the results were discussed in detail below.

3.1 Physical Properties

The highest and the lowest values across the depth were discussed below in detail. Bulk densities (BD) of the samples analyzed varied from 1.55 to 1.31 Mg m⁻³ at three different depths and are significantly different. Among all, the treatment receiving FYM @ 12.5 t ha⁻¹ have the lowest bulk density (1.31 Mg m⁻³) at 0-15 cm when compared to control (1.55 Mg m⁻³) at 30-45 cm depth. The values were observed to increase with the increase in depth. Maximum reduction in bulk density was recorded when FYM was applied along with chemical fertilizers which may be due to the positive effect of organic manure on soil aggregation. The higher bulk density in control and in only N fertilizer treated plots may be due to low organic matter content in soil and formation of compact layer [17]. However, porosity varied along with the depth with a decreasing trend and was significantly different. The values ranged from 49.02% (FYM @ 12.5 t ha⁻¹) at 0-15 cm depth to 23.26% (control) at 30-45 cm depth which may be due to the positive effect of organic manure on soil aggregation [18]. A similar trend was recorded in field capacity (33.21% in FYM @ 12.5 t ha⁻¹ at 0-15 cm to 20.71% in control at 30-45 cm depth). Permanent Wilting Point was seen highest in FYM @ 12.5 t ha⁻¹ but not in a decreased trend and the values ranged from 18.80% (FYM @ 12.5 t ha⁻¹ at 30-45 cm) to 13.89% (control at 0-15 cm depth). Available Water was observed to follow the same

trend of porosity and field capacity (15.54% in FYM @ 12.5 t ha⁻¹ at 0-15 cm to 8.30 % in control at 30-45 cm depth) which decreased with depth and are significantly different. Hydraulic conductivity ranged from 3.11cm hr⁻¹ (FYM @ 12.5 t ha⁻¹) at 0-15 cm to 1.80 cm hr⁻¹ (control) at 30-45 cm depth, the decrease in SHC values at lower depth may also be due to an increase in the clay content of soil [19]. Clay offers a higher resistance to the movement of water because of its high proportion of micropores that store water in film or gyroscopically.

To conclude, the soil physical properties were well maintained under organics alone which was almost similar with organics applied along with inorganics. It was also observed that the same trend was seen with all three depths with a decreasing trend with increasing depth.

3.2 Chemical Properties

Soil reaction (pH) varied from slightly alkaline to slightly neutral with an increase in depth where the highest pH was observed in Control (8.16) at 0-15 cm depth and the lowest in the treatment FYM @ 12.5 t ha⁻¹ (7.60) at 30-45 cm depth may probably be due to organic acids released during the decomposition of organic matter resulting lower pH Liang et al. [20], Arulmozhiselvan et al. [21] and Malarkodi et al. [22]. Correspondingly Electrical Conductivity (EC) ranged from 0.19 dS m⁻¹ in FYM @ 12.5 t ha⁻¹ at 0-15 cm to 0.10 dS m⁻¹ in control at 30-45 cm depth. Organic Carbon was significantly different in between the treatments and ranged from 3.60 g kg⁻¹ (FYM @ 12.5 t ha⁻¹) at 0-15 cm to 0.78 g kg⁻¹ (Control) at 30-45 cm depth which was in a decreasing trend. A decreasing trend with an increase in depth was followed for all the available nutrients with,

Table 5. Effect of continuous application of fertilizers or manures on physical properties of soil under different depths

	Bulk Density (Mg m ⁻³)			Porosity (%)			Field Capacity (%)		
	0-15	15-30	30-45	0-15	15-30	30-45	0-15	15-30	30-45
Control	1.50	1.53	1.55	24.81	23.66	23.26	27.16	23.98	20.71
100 % RDF	1.43	1.47	1.49	32.14	30.88	30.60	30.42	26.25	21.18
50% RDF	1.48	1.52	1.54	27.41	26.52	26.15	29.15	24.66	20.80
50% N (Crop residues)	1.40	1.44	1.46	37.06	36.69	36.50	30.10	24.77	21.09
50 % N (FYM)	1.38	1.41	1.43	37.93	37.32	37.14	30.87	25.99	21.14
50 % Inorganic N+ 50% organic N (crop residues) + P (50%) + K (50%)	1.35	1.40	1.42	42.57	41.26	41.13	31.63	27.43	25.34
50 % Inorganic N+ 50% organic N (FYM) + P (50%) + K (50%)	1.33	1.38	1.40	46.00	44.83	44.76	31.92	27.36	22.01
100 % RDF + 25 kg ZnSO ₄ ha ⁻¹	1.43	1.45	1.47	33.57	33.33	33.09	31.00	26.61	21.64
FYM - 12.5 t ha ⁻¹	1.31	1.35	1.37	49.02	47.97	47.95	33.21	29.25	27.10
Mean	1.40	1.44	1.46	36.72	35.83	35.62	30.61	26.26	22.33
CD	0.051	0.052	0.053	2.900	2.886	2.413	1.454	1.248	2.570
SE(d)	0.024	0.024	0.025	1.368	1.361	1.138	0.686	0.588	1.213

Table 6. Effect of continuous application of fertilizers or manures on physical properties of soil under different depths

	Permanent Wilting Point (%)			Available Water Content (%)			Hydraulic Conductivity (cm hr ⁻¹)		
	0-15	15-30	30-45	0-15	15-30	30-45	0-15	15-30	30-45
Control	13.89	13.89	15.20	13.27	10.09	5.51	2.51	2.07	1.80
100 % RDF	16.24	15.96	14.68	14.18	10.29	6.50	2.68	2.29	1.95
50% RDF	15.06	15.00	15.13	14.09	9.66	5.67	2.57	2.24	1.91
50% N (Crop residues)	15.84	14.14	14.36	14.26	10.63	6.73	2.62	2.40	2.02
50 % N (FYM)	16.65	15.20	14.20	14.22	10.79	6.94	2.76	2.46	2.09
50 % Inorganic N+ 50% organic N (crop residues) + P (50%) + K (50%)	17.22	16.50	17.41	14.41	10.93	7.93	2.81	2.52	2.21
50 % Inorganic N+ 50% organic N (FYM) + P (50%) + K (50%)	17.52	16.19	13.76	14.40	11.17	8.25	2.98	2.53	2.27
100 % RDF + 25 kg ZnSO ₄ ha ⁻¹	16.80	16.22	14.96	14.20	10.39	6.68	2.78	2.31	2.01
FYM - 12.5 t ha ⁻¹	17.67	17.97	18.80	15.54	11.28	8.30	3.11	2.65	2.32
Mean	16.32	15.67	15.39	14.29	10.58	6.95	2.76	2.39	2.06
CD	0.777	0.575	0.565	1.144	1.342	1.134	0.446	0.385	0.334
SE(d)	0.366	0.271	0.266	0.539	0.633	0.535	0.210	0.182	0.158

Table 7. Effect of continuous application of fertilizers or manures on chemical properties of soil under different depths

	pH			Electrical Conductivity (dS m ⁻¹)			Organic Carbon (g kg ⁻¹)		
	0-15	15-30	30-45	0-15	15-30	30-45	0-15	15-30	30-45
Control	8.16	8.09	7.89	0.12	0.10	0.10	1.40	1.10	0.78
100 % RDF	8.03	7.84	7.71	0.12	0.1	0.1	2.30	1.90	1.80
50% RDF	8.05	7.95	7.86	0.13	0.11	0.11	1.80	1.50	1.20
50% N (Crop residues)	7.94	7.91	7.78	0.13	0.11	0.11	2.00	1.80	1.50
50 % N (FYM)	7.91	7.87	7.73	0.13	0.12	0.12	2.20	1.90	1.50
50 % Inorganic N+ 50% organic N (crop residues) + P (50%) + K (50%)	7.85	7.73	7.62	0.15	0.13	0.13	3.10	2.40	2.30
50 % Inorganic N+ 50% organic N (FYM) + P (50%) + K (50%)	7.84	7.71	7.65	0.16	0.15	0.14	3.40	2.50	2.50
100 % RDF + 25 kg ZnSO ₄ ha ⁻¹	7.89	7.81	7.68	0.14	0.13	0.13	2.40	2.10	2.00
FYM - 12.5 t ha ⁻¹	7.81	7.63	7.60	0.19	0.18	0.16	3.60	3.20	2.80
Mean	7.94	7.84	7.72	0.14	0.13	0.12	2.47	2.04	1.82
CD	0.777	0.575	0.565	1.144	1.342	1.134	0.416	0.344	0.312
SE(d)	0.366	0.271	0.266	0.539	0.633	0.535	0.196	0.162	0.147

Table 8. Effect of continuous application of fertilizers or manures on Chemical properties of soil under different depths

	Nitrogen (kg ha ⁻¹)			Phosphorus (kg ha ⁻¹)			Potassium (kg ha ⁻¹)		
	0-15	15-30	30-45	0-15	15-30	30-45	0-15	15-30	30-45
Control	100	90	69	10.86	8.60	4.53	308	215	137
100 % RDF	137	112	94	13.58	9.05	7.24	372	295	218
50% RDF	125	103	85	11.09	8.60	7.01	313	221	159
50% N (Crop residues)	131	106	88	11.31	8.70	7.24	329	252	174
50 % N (FYM)	134	109	91	11.31	8.82	7.24	356	256	201
50 % Inorganic N+ 50% organic N (crop residues) + P (50%) + K (50%)	144	120	100	14.03	9.38	7.69	395	321	240
50 % Inorganic N+ 50% organic N (FYM) + P (50%) + K (50%)	147	128	107	14.48	9.41	8.01	404	326	246
100 % RDF + 25 kg ZnSO ₄ ha ⁻¹	141	116	97	13.58	9.13	7.47	381	304	227
FYM - 12.5 t ha ⁻¹	150	137	110	20.59	9.73	8.37	416	339	255
Mean	134	113	93.4	13.4	9.05	7.20	364	281	206
CD	10.80	9.135	7.530	2.215	1.460	1.176	13.38	10.42	7.69
SE(d)	5.095	4.309	3.552	1.045	0.689	0.555	6.31	4.92	3.63

Nitrogen (N) ranging from 150 kg ha⁻¹ to 69 Kg ha⁻¹, Phosphorus (P) from 20.59 kg ha⁻¹ to 4.53 kg ha⁻¹, Potassium from 416 kg ha⁻¹ to 137 kg ha⁻¹. This might be due to integrated nutrient application, higher microbial population, and high organic carbon, organic form of nutrients are converted to inorganic [23].

To summarize, the treatment which received organics alone was noticed to perform well in maintaining soil chemical properties which were nearly similar to with the treatment of organics applied along with inorganics. The values were observed to decrease with increase with depth with the same trend.

4. CONCLUSION

The present investigation revealed that the application of organics viz., T₉- FYM @ 12.5 t ha⁻¹ resulted in improving the soil's physical, chemical properties. Improving these properties may directly improve soil health which increases the production and productivity of crops. The organics was then followed by application of organics along with inorganics viz., T₇- 50 % Inorganic N+ 50% organic N (FYM) + P (50%) + K (50%) and T₆- 50 % Inorganic N+ 50% organic N (crop residues) + P (50%) + K (50%). It was also observed that all the values were decreased with an increase in depth. So it is concluded that the application of organics is best in improving soil health, if not available it can be substituted with organics+ inorganics.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCE

1. Perrott KW, Sarathchandra SU, Dow BW. Seasonal and fertilizer effects on the organic cycle and microbial biomass in a hill country soil under pasture. *Australian Journal of Soil Research*. 1992;30:383-394.
2. Steinshamn H, Thuen E, Bleken MA, Brenoe UT, Ekerholt G and Yri G. Utilization of nitrogen (N) and phosphorus (P) in an organic dairy farming system in Norway. *Agriculture Ecosystems and Environment*. 2004;104:509-522.
3. Avnimelech Y. Organic residues in modern agriculture. In *The role of organic manure in modern agriculture*, ed. Y. Chen and Y. Avnimelech, 1-9. Dordrecht, the Netherlands: Martinus Nyhoff; 1986.
4. Singh R and Rai RK. Yield attributes, yield, and quality of soybean (*Glycine max*) as influenced by integrated nutrient management. *Indian Journal of Agronomy*. 2004;49(4):271-274.
5. Bhattacharyya R, Kundu S, Prakash R and Gupta HS. Sustainability under combined application of mineral and organic fertilizers in a rainfed soybean of the Indian Himalayas. *European Journal of Agronomy*. 2008;28(1):33-46.
6. Wu SC, Cao ZH, Li ZG, Cheung KC and Wong MH. Effects of biofertilizer containing N-fixer, P and K solubilizers and AM fungi on maize growth: A greenhouse trial. *Geoderma*. 2005;125: 155-166.
7. Singh AK. Soil quality parameters as influenced by management practices in rice-wheat and maize-wheat cropping systems. In *Proceedings of the 19th World Congress of Soil Science: Soil solutions for a changing world*, Brisbane, Australia, 1-6 August 2010. Symposium 3.3.1 Integrated Nutrient Management. 2010;278-281.
8. More SD and Hangarge DS. Effect of integrated nutrient supply on crop productivity and soil characteristics with cotton– sorghum cropping sequence in Vertisol. *Journal of Maharashtra Agricultural University*. 2003;28(1):8-12.
9. Gupta RP, Dakshinamoorthy C. *Procedures for physical analysis of soil and collection of agrometeorological data*. New Delhi: Indian Agricultural Research Institute. 1980;293.
10. Richards LA. *Diagnosis and Improvement of Saline and Alkali Soils*. Handbook 60. Soil Tillage Res. 1954;112:167–74.
11. Jackson ML. *Soil chemical analysis*. Prentice-Hall of India Pvt. Ltd. New Delhi, India: 1973a; 327-350.
12. Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil sci*. 1934; 37(1), pp.29-38.
13. Subbiah BV, Asija GL. Alkaline method for determination of mineralizable nitrogen. *Curr. Sci*. 1956; 25 (2):259-260.
14. Olsen, SR. Estimation of available phosphorus in soils by extraction with sodium bicarbonate, United States Department Of Agriculture; Washington; 1954.

15. Stanford G, English L. "Use of the flame photometer in rapid soil tests for K and Ca." *Agronomy Journal*.1949;41(9):446-447.
16. Gomez K A, et al. *Statistical procedures for agricultural research*, John Wiley & Sons; 1984.
17. Islam MM, Karim AJMS, Jahiruddin M, Majid NM, Miah MG, Ahmed MM and Hakim MA. Effects of organic manure and chemical fertilizers on crops in the radish-stem amaranth Indian spinach cropping pattern in homestead area. *Australian Journal of Crop Science*. 2011;5:1370-1378
18. Meng L, Ding W, Cai Z. Long-term application of organic manure and nitrogen fertilizer on N₂O emissions, soil quality and crop production in a sandy loam soil. *Soil Biol. and Biochem*. 2005;37(11):2037-2045.
Available:<https://doi.org/10.1016/j.soilbio.2005.03.007>.
19. Edoga RN. Comparison of saturated hydraulic conductivity measurement methods for samaru-nigeria soils. *Libyan Agriculture Research Center Journal International*. 2010;1:269-273.
20. Liang Q, Chen H, Gong Y, Fan M, Yang H, Lal R, and Kuzyakov Y. Effects of 15 years of manure and inorganic fertilizers on soil organic carbon fractions in a wheat-maize system in the North China Plain. *Nutrient cycling in agroecosystems* 2012;92(1):21-33.
21. Arulmozhiselvan K, Sathya S, Elayarajan M, and Malarkodi M. Soil fertility changes and crop productivity of finger millet under continuous fertilization and manuring in finger millet-maize cropping sequence. *Res. Environ. Life Sci*. 2015;8(4):751-756.
22. Malarkodi M, Elayarajan M, Arulmozhiselvan K, and Gokila B. Long-term impact of fertilizers and manures on crop productivity and soil fertility in an alfisol. *The Pharma Innovation*. 2019;8(7):252-256.
23. Dhaliwal, SS, Naresh RK, Mandal A, Walia MK, Gupta RK, Singh R, and Dhaliwal MK. Effect of manures and fertilizers on soil physical properties, build-up of macro and micronutrients and uptake in soil under different cropping systems: a review. *Journal of Plant Nutrition*. 2019;42(20):2873-2900.

© 2021 Venkatesh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/79880>