



Modulatory Effect of Aqueous Ginger (*Zingiber officinale*) Extract on Gentamicin Sulfate Induced Oxidative Stress and Nephrotoxicity in Female Albino Rats

Henry Wealth Oyarieme ^a, Akinniyi Adediran Osuntoki ^a,
Holiness S.A. Olasore ^a, Ogheneovo Obros ^a,
Jeromi Otiti ^b and Zakariyah Abdullahi Abiodun ^{c*}

^a Department of Biochemistry, Faculty of Basic Medical Sciences, College of Medicine, University of Lagos, Idi - Araba, Lagos, Nigeria.

^b Department of Zoology and Environmental Biology, Faculty of Science, Olabisi Onabanjo University, Ago-Iwoye, Ogun State, Nigeria.

^c Chemical Engineering Department, Federal University of Technology, Minna, P.M.B. 65, Niger State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajbgmb/2024/v16i9404>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121844>

Original Research Article

Received: 24/06/2024
Accepted: 27/08/2024
Published: 30/08/2024

*Corresponding author: E-mail: Zakariyahabdullahi20@gmail.com; Zabiodun31@gmail.com;

Cite as: Oyarieme, Henry Wealth, Akinniyi Adediran Osuntoki, Holiness S.A. Olasore, Ogheneovo Obros, Jeromi Otiti, and Zakariyah Abdullahi Abiodun. 2024. "Modulatory Effect of Aqueous Ginger (*Zingiber officinale*) Extract on Gentamicin Sulfate Induced Oxidative Stress and Nephrotoxicity in Female Albino Rats". *Asian Journal of Biochemistry, Genetics and Molecular Biology* 16 (9):40-58. <https://doi.org/10.9734/ajbgmb/2024/v16i9404>.

ABSTRACT

Acute renal failure is a common complication of gentamicin, an aminoglycoside antibiotic widely used against gram-negative infections, often mediated by oxidative stress. This study investigates the potential protective effects of aqueous ginger extract against gentamicin-induced nephrotoxicity in female albino rats. Rats were divided into four groups: control (distilled water and feed), ginger extract (2.14 ml/kg), gentamicin (100 mg/kg), and combined ginger extract with gentamicin. The combination of ginger extract with gentamicin has garnered attention in recent research due to its potential protective effects against the nephrotoxic side effects of gentamicin, an aminoglycoside antibiotic commonly used to treat bacterial infections. Biochemical assays measured blood urea nitrogen (BUN), creatinine, cholesterol, triglycerides, albumin, white blood cell count, and packed cell volume. Kidney homogenates were analyzed for catalase (CAT), superoxide dismutase (SOD), reduced glutathione (GSH), glutathione transferase (GST) activities, and malondialdehyde (MDA) levels. Histopathological evaluations of kidney tissues were also performed. Gentamicin administration significantly increased BUN and creatinine levels, decreased GSH levels, and increased MDA levels, indicating severe nephrotoxicity and oxidative stress. Ginger extract administration did not alleviate gentamicin-induced renal damage or oxidative stress, despite reducing serum cholesterol levels. The study also observed a reduction in CAT activity with ginger extract administration. Histopathological analysis revealed clear nephritis in the gentamicin-treated groups, particularly exacerbated in the group receiving both gentamicin and ginger extract. These outcomes suggest that while aqueous ginger extract did not mitigate gentamicin-induced renal injury in this study, its cholesterol-lowering effects highlight potential benefits in managing atherosclerosis. Further research is necessary to explore under varying conditions which ginger extract may discuss renal protection.

Keywords: Ginger; glutathione transferase (GST); albumin; histopathological analysis;therosclerosis.

1. INTRODUCTION

“Nephrology is the branch of medicine that focuses on the kidneys and kidney-related diseases. The kidneys are vital organs that play a crucial role in maintaining the body's homeostasis by filtering waste products and excess fluids from the bloodstream and excreting them as urine. This process is essential for the removal of harmful substances and the regulation of various physiological functions. In both humans and dogs, the kidneys are situated in the rear portion of the abdomen. Positioned above each kidney is the adrenal gland (also known as the suprarenal gland), and each kidney is enveloped by two layers of fat (the perirenal and pararenal fat), which serve to cushion it” [1]. “The fundamental functional unit of the kidney is the nephron, of which there are more than a million within the cortex and medulla of each normal adult kidney. It regulates water and soluble substances (especially electrolytes) in the body by first filtering the blood under pressure, then reabsorbing some necessary fluid and molecules back into the blood while eliminating those that are not needed. Reabsorption and secretion are accomplished with other co-transport and counter-transport mechanisms established in the nephrons and are associated with collecting ducts. This segment of

the nephron is crucial to the process of water conservation by the organism. In the presence of antidiuretic hormone (ADH; also called vasopressin), these ducts become less permeable to water, facilitating its reabsorption, thus concentrating the urine and reducing its volume. When the organism must eliminate water, such as after excess fluid intake, the reduction of ADH is decreased, and the collecting tubule becomes less permeable to water, rendering the urine dilute and abundant. Failure of the organism to decrease ADH production appropriately, a condition known as syndrome of inappropriate ADH (SIADH), may lead to water retention and dangerous dilution of the body fluid, which may cause severe neurological damage. Failure to produce ADH (or inability of the collecting ducts to respond to it) may cause excessive urination, called diabetes insipidus (DI). The functions of the kidney include excretion of a variety of waste products produced by metabolism, including the nitrogenous wastes, urea (from protein catabolism) and uric acid (from nucleic acid metabolism) and water. Homeostatic functions such as acid-base balance, regulation of electrolyte concentration, control of blood volume, and regulation of blood pressure. Plasma Volume, Any significant rise or drop in plasma osmolality is detected by the hypothalamus, which communicates directly with

the posterior pituitary gland. A rise in osmolality causes the gland to secrete antidiuretic hormone, resulting in water reabsorption by the kidney and an increase in urine concentration. The two factors work together to return the plasma osmolality to its normal levels. The kidneys also secrete a variety of hormones, including erythropoietin, urodilatin, renin, and vitamin D. Kidney diseases and disorders can be either congenital or acquired. The congenital conditions include congenital hydronephrosis, congenital obstruction of the urinary tract, duplicated ureter, horseshoe kidney, polycystic kidney disease, renal dysplasia, unilateral small kidney, and medullary sponge kidney. Acquired conditions include diabetic nephropathy, glomerulonephritis, hydronephrosis, interstitial nephritis, kidney tumors (Wilms tumor, renal cell carcinoma), lupus nephritis, minimal change disease, pyelonephritis, acute or chronic renal failure, and kidney stones. Renal calculi, also known as kidney stones, are solid concretions (crystal aggregations) of dissolved minerals in urine, which typically form inside the kidneys or ureters. The term nephrolithiasis refers to the presence of calculi in the kidneys and urinary tracts, respectively" [2]. "Kidney stones can be attributed to underlying metabolic conditions, such as renal tubular acidosis, Dent's disease, and medullary sponge kidney, and many healthcare facilities will screen for such disorders in patients with recurrent kidney stones, although most stones arise spontaneously. Symptoms of kidney stones include colicky pain radiating from the flank to the groin, hematuria due to damage to the wall of the ureter or urethra, dysuria when passing stones, oliguria due to obstruction of the bladder or urethra by a stone, or extremely rarely, simultaneous obstruction of both ureters by a stone, and nausea or vomiting due to an embryological link with the intestine. Preventive strategies involve dietary modification and sometimes drug therapy, with the goal of reducing the excretory load on the kidneys" [3]. "Ginger (*Zingiber officinale*) is a versatile plant with a long history of traditional medicinal use. Its rhizome has been employed to alleviate various ailments, including joint inflammation, cold symptoms, sore throats, fever, cramps, and constipation" [4]. "Additionally, ginger exhibits antioxidant properties and cytoprotective effects, making it a potential treatment for microbial infections, inflammatory conditions, and emesis" [5]. "The bioactive constituents of *Zingiber officinale*, such as zingerone, gingerdiol, zingibrene, gingerols, and shogaols, are responsible for its antioxidant activities" [6].

Interestingly, ginger has been found to play a preventive role in ischemia/reperfusion (I/R) injury in rat kidneys, suggesting its potential in protecting against kidney damage caused by reduced blood flow and subsequent restoration [7]. Furthermore, ginger has demonstrated an inhibitory effect on cisplatin-induced nephrotoxicity in rats, indicating its possible therapeutic application in mitigating drug-induced kidney injury [7]. While ginger is commonly used for managing morning sickness during the first trimester of pregnancy, recent literature reviews have concluded that its safety during pregnancy has not been definitively established [8]. Ginger rhizomes are also extensively utilized in foods, especially in Asian cuisines, due to their medicinal benefits. The primary bioactive constituents in *Zingiber officinale* (ginger) are the pungent vanilloids, 6-gingerol and 6-paradol, along with other phenolic compounds such as shogaols and zingerone. Ginger rhizomes are also rich in essential minerals, including high amounts of iron (54-62 mg/100g) and calcium (1.0%-1.5%). The antioxidant, antitumor, and anti-inflammatory properties of ginger are primarily attributed to its pungent constituents, particularly 6-gingerol. The antioxidant effects of ginger have been demonstrated in various studies, including one where ginger was shown to overcome the oxidative stress induced by the pesticide malathion in rats [9]. Ginger is also recognized for its potent antitumor activities, as evidenced by in vitro experiments, with the vanilloids 6-gingerol and 6-paradol playing a key role in inhibiting cellular proliferation and inducing apoptosis in cancer cells [10]. Furthermore, the anti-inflammatory effects of 6-gingerol have been studied in mouse models, where it was found to suppress the inflammatory response induced by the phorbol ester TPA [11]. Ginger contains diverse bioactive compounds, such as gingerols, shogaols, and paradols, and possesses antioxidant and anti-inflammatory properties that might help reduce the levels of inflammation and oxidative stress in neurodegenerative diseases. Several inflammatory, oxidative, and immunopathological parameters are involved in their pathogenesis, and drugs used for treatment are of limited efficacy and can also generate adverse side effects. It seems that ginger, given its antioxidant, immunomodulatory, and anti-inflammatory capacity, has the ability to intercept the main elements involved in the development of multiple sclerosis as well as to attenuate the symptoms of neurological diseases including Parkinson's, Alzheimer's, migraine, and epilepsy [12]. Similarly, ginger oil has been shown to

significantly reduce joint swelling in a rat model of severe arthritis, with the oleoresin constituents and phenolic compounds, such as paradol and shogaol, exhibiting inhibitory effects on the cyclooxygenase-2 (COX-2) enzyme, a crucial mechanism in the regulation of the inflammatory process [13]. These findings suggest that ginger and its bioactive compounds may have therapeutic potential in the management of various inflammatory conditions. The health benefits of ginger are mainly attributed to its phenolic compounds, such as gingerols and shogaols. Accumulated investigations have demonstrated that ginger possesses multiple biological activities, including antioxidant, anti-inflammatory, antimicrobial, anticancer, neuroprotective, cardiovascular protective, respiratory protective, antiobesity, antidiabetic, anti-nausea, and antiemetic activities [10]. Inclusion of ginger or ginger extracts in nutraceutical formulations could provide valuable protection against neurodegenerative diseases.

2. MATERIALS AND METHODS

2.1 Sample Preparation

Fresh samples of ginger (*Zingiber officinale*), belonging to the Zingiberaceae family, were procured from Mushin Market, Mushin, Lagos State, Nigeria. Prior to extraction, the rhizomes were thoroughly washed to remove any surface contaminants. The scaly skin was then peeled off, and the rhizomes were re-washed with distilled water to eliminate unwanted materials such as sand, debris, and other contaminants. This initial cleaning process is crucial for ensuring the purity of the extract, as impurities can interfere with subsequent analyses and applications [14]. After cleaning, the peeled ginger rhizomes were reduced to smaller particles by grating. The grated ginger was then subjected to hot water extraction, where it was boiled for 10 minutes using a ratio of 50g of ginger to 200ml of distilled water. This method of extraction is effective for obtaining bioactive compounds, as heat can facilitate the release of essential oils and other soluble constituents from the ginger rhizome [15,16]. Following extraction, the solution was filtered using glass wool to separate the solid residues from the liquid extract. The resulting extract was then concentrated by applying direct heat, which helps to evaporate excess water and enhance the concentration of active ingredients. Finally, the concentrated extract was dried in an oven at 80°C to yield a dry product. The process resulted in 10.5g of dried extract per 50g of wet ginger

rhizome, indicating a substantial yield of bioactive compounds.

2.2 Experimental Design

Twenty-nine female albino rats weighing between 100-180 g were procured from the University of Ibadan for the purpose of this research. Based on their intended uses, the animals were redistributed into four different groups (Groups A, B, C, and D) and housed in well-ventilated iron cages at room temperature. They were fed a standard rat diet and provided with drinking water. The room and the cages were regularly cleaned and disinfected to avoid infections. The acclimatization period lasted for two weeks, during which all the groups were fed the rat feed and had access to drinking water throughout the day.

Group A: Received rat feed and distilled water.

Group B: Received rat feed, distilled water, and ginger aqueous extract (2.14 mL/kg or 146 mg/kg).

Group C: Received rat feed, distilled water, and gentamicin (100 mg/kg) without ginger extract.

Group D: Received rat feed, distilled water, ginger extract (2.14 mL/kg), and gentamicin (100 g/kg).

2.3 Blood Collection for Hematological and Biochemical Analysis

Blood collection was performed under light anesthesia using ketamine hydrochloride at a dosage of 5 mg/kg body weight. This level of sedation was chosen to minimize stress and discomfort for the subjects during the procedure, ensuring a humane approach to blood sampling [17]. Blood was obtained via cardiac puncture, a technique that allows for the collection of a sufficient volume of blood while minimizing trauma to the animal [18]. The serum was prepared from the obtained blood samples by centrifugation at 3000 rpm for 20 minutes. This centrifugation speed and duration are optimal for separating serum from cellular components, ensuring the clarity and integrity of the serum for subsequent biochemical assays [19]. The serum was then subjected to various biochemical assays, including measurements of blood urea nitrogen (BUN), creatinine, cholesterol, triglycerides, and albumin. These parameters are essential for assessing renal function, lipid metabolism, and overall health status [20]. A portion of the blood was collected in a heparinized bottle for white blood cell count and packed cell volume.




Fig. 1. Female albino rats weighing between 100-180 g

2.4 Kidney Homogenate Preparation

From the sacrificed rats, the kidneys were dissected out, blotted on filter paper, washed in 0.9% saline, and weighed immediately. Kidney homogenate was prepared by homogenizing 0.5 g of kidney tissue in phosphate buffer. The prepared homogenate was centrifuged at 3000 g for 20 minutes, and the supernatant was collected and stored in a refrigerator for further use, including assays for catalase, superoxide dismutase, lipid peroxidation (thiobarbituric acid), reduced glutathione (GSH), and glutathione-S-transferase, which were performed within two weeks.

2.5 Histopathological Evaluation

A kidney was taken from a rat (two from each group) for histopathological evaluation. The histopathological evaluation involved collecting kidney samples from four groups of rats. Group A (GA) served as the control, receiving distilled water and standard rat feed. Group B (GB) was given ginger extract and rat feed. Group C (GC) received rat feed and an overdose of gentamicin, and Group D (GD) was administered feed, ginger extract, and an excessive dose of gentamicin. The collected samples were fixed in 10% formalin, processed, and sectioned at 4 microns. The tissue sections were then stained using the conventional hematoxylin and eosin method, and the slides were thoroughly interpreted and summarized according to the respective groups.

2.6 Hematological Analysis

2.6.1 Determination of packed cell volume (PCV) [Hematocrit]

The rat blood collected in a heparinized container was drawn into a hematocrit tube, sealed at one end with Plasticine, and centrifuged for 20 minutes at 3000 rpm. A hematocrit reader was used to determine the volume of the red cells relative to the whole column, which is known as the hematocrit reading or packed cell volume.

2.6.2 White blood cell count

The method described by [9] was used. The rat blood in the heparinized bottle was drawn to the 0.5 mark on the stem of a white cell pipette and diluted with a diluting fluid to the 11 mark immediately above the bulb. Using an improved Neubauer Chamber, the cells in the four corner squares and those in the central square millimeter of the ruled area were counted.

2.6.3 Biochemical analysis

Catalase activity was assessed using the method by [10], measured spectrophotometrically by the decrease in absorbance at 240 nm (A_{240}) as hydrogen peroxide (H_2O_2) decomposed, with a Beckman spectrophotometer (USA) and a molar extinction coefficient of $40.0 \text{ m}^{-1} \text{ cm}^{-1}$. The materials used included a phosphate buffer (50 mM, pH 7.0) prepared from KH_2PO_4 and $Na_2HPO_4 \cdot 2H_2O$, and hydrogen peroxide (30

mM). In the procedure, 0.2 mL of the sample was mixed with 1.8 mL of 30 mM H₂O₂, or 0.1 mL of kidney homogenate supernatant, with phosphate buffer as the blank. Absorbance was recorded at 240 nm every 30 seconds for 1 minute.

2.6.4 Superoxide Dismutase (SOD) Assay

Superoxide dismutase (SOD) activity was assayed according to the method of [11]. 3 mL of 50 mM Na₂CO₃ buffer + 0.02 mL (20 µL) of the sample (kidney homogenate supernatant) + 0.03 mL (30 µL) of epinephrine, and the absorbance was measured at 480 nm for 3-5 minutes. The blank used was 3 mL of buffer + 0.02 mL of water + 0.03 mL of the substrate (sample).

$$\text{Enzyme Activity} = \frac{\frac{\Delta OD}{\min 480} * V_t}{\sum x * V_s} = \text{Amount of O}_2, \text{ produced per ml}$$

Where:

V_t=Total volume of reaching sample

V_s=Volume of sample

∑ x= 4020.0m⁻¹ cm⁻¹

The superoxide anion radical (O₂⁻) spontaneously dismutates to O₂ and H₂O₂ rapidly. (Ω 105 m⁻¹ 5⁻¹ at PH7). SOD catalyses the dismutation of superoxide into oxygen and hydrogen peroxide.

2.6.5 Thiobarbituric acid assay (TBARS)

The method according to [12] was used. Malondialdehyde, formed from the breakdown of polyunsaturated fatty acids, serves as a convenient index for determining the extent of the peroxidation reaction. 1.0 mL of the kidney homogenate supernatant was mixed with 2.0 mL of the TCA-TBA-HCl reagent and thoroughly mixed. The solution was heated for 15 minutes in a boiling water bath. After cooling, the flocculent precipitate was removed by centrifugation at 1000 rpm for 10 minutes. The absorbance of the sample was determined at 535 nm against a blank that contained all the reagents minus the kidney homogenate supernatant. The malondialdehyde concentration in the sample was calculated using an extinction coefficient of 1.56 × 10⁵ m⁻¹ cm⁻¹ [13].

$$\text{Malondialdehyde (nmol)} = \frac{OD * V_t * F}{\sum V_s}$$

Where:

OD= Optical density (absorbance reading)

V_t= Total volume of reaction mixture

V= Volume of the sample

E = Standard coefficient (1.56 x10⁵m⁻¹ cm⁻¹)

F = dilution factor (Optional)

2.6.6 Glutathione-S-transferase (GST) assay

The method of Habig (1974) was used. The reaction was measured by observing the conjugation of 1-chloro-2,4-dinitrobenzene (CDNB) with reduced glutathione (GSH). This was done by monitoring the increase in absorbance at 340 nm. One unit of the enzyme would conjugate 10 nanomoles of CDNB with reduced glutathione per minute at 25°C [15]. Glutathione-S-Transferase catalyzed the reaction (binding) of GSH (reduced glutathione) and CDNB. 1 mL of phosphate buffer (0.2 M phosphate buffer, pH 6.5), 0.1 mL of CDNB (20 mM CDNB) and 1.7 mL of distilled water were added, and the mixture was incubated at 37°C for 5 minutes. Then, 0.1 mL of the kidney supernatant homogenate and 0.1 mL (20 mM GSH) were added, and the absorbance was taken for 5 minutes at 340 nm.

2.6.7 Reduced glutathione (GSH) assay

The procedure of [16] was used for this analysis. GSH is known as a substrate in both conjugation and reduction reactions. The level of GSH was determined spectrophotometrically. 100 µL of the kidney supernatant homogenate was added to 1 mL of 0.2 M Tris-EDTA buffer (pH 8.2), 0.9 mL of 20 mM EDTA (pH 4.7), and 20 µL of 10 mM DTNB. After 30 minutes of incubation at room temperature, the mixture was centrifuged, and the absorbance of the supernatant was read against distilled water at 412 nm.

$$\text{GSH} = \frac{OD * V_t}{\sum V_s}$$

Where:

OD = Absorbance

V_t = Total volume of reaction mixture

V_s= Volume of sample in reaction mixture

∑ = Extinction coefficient =13,600mcm.

2.6.8 Triglyceride assay

Serum triglycerides were analyzed using a standard enzymatic procedure from the Hirnen Gesellschaft for Biochemical and Diagnostic mbH (Germany). The triglycerides were quantified after enzymatic hydrolysis with lipases. The indicator is quinoneimine, which is formed from hydrogen peroxide, 4-amino-antipyrine, and 4-chlorophenol under the catalytic influence of

peroxidase. The procedure was outlined by [17]. The absorbance of the sample (AA sample) and the standard ΔA (STD) against the reagent blank at 500nm was taken within 60 minutes. The Triglycerides Concentration (C) was calculated using.

$$C = \frac{2.28 * \Delta A \text{ sample } (\frac{mmol}{L})}{\Delta A(STD)}$$

2.6.9 Total cholesterol assay

The serum total cholesterol was analyzed using standard enzymatic procedures (Human Biochemical, Germany) outlined by [18]. The assay principle involved the determination of cholesterol after enzymatic hydrolysis and oxidation. For the assay, the following reagents were mixed: phosphate buffer (100 mmol/L, pH 6.5), 4-amino phenazone (0.3 mmol/L), phenol (5 mmol/L), peroxidase (>5 ku/L), cholesterol oxidase (>100 u/L), and sodium oxide (0.05%). The standard used was 3 mL of cholesterol at 5.17 mmol/L. In the pipetting scheme, 10 μ L of the sample or standard was added to 1000 μ L of reagent (RGT), while the reagent blank consisted of 1000 μ L of reagent alone. The mixture was incubated at 37°C, and absorbance was measured at 500 nm within 60 minutes, comparing the sample or standard against the reagent blank. The serum total cholesterol was calculated as follows

$$\text{Total serum cholesterol} = \frac{5.17 * \Delta A \text{ sample}}{\Delta A (\text{Standard})} (\text{mmol/L})$$

2.6.10 Albumin assay

Serum albumin was analyzed using a standard procedure from Human Gesellschaft for Biochemical and Diagnostic mbH (Germany) involving a colorimetric method, as outlined by [19]. In this method, albumin is determined after it forms a colored complex with bromocresol green in a citrate buffer. For the assay, the following reagents were used: citrate buffer (30 mmol/L), bromocresol green (260 μ mol/L), albumin standard (4 g/dL or 40 g/L), and sodium azide (0.05%). In the pipetting scheme, 10 μ L of the sample or standard was added to 1000 μ L of reagent (RGT), while the reagent blank consisted of 1000 μ L of reagent alone. The absorbance of the sample and the standard was measured against the reagent blank at 578 nm within 30 minutes (ΔA). Albumin Concentration was calculated using:

$$C = \frac{40 * \Delta A \text{ sample}}{\Delta A (\text{Standard})} (\text{g/L})$$

2.6.11 Test for creatinine

Serum creatinine was analyzed using the standard procedure from Randox Laboratory Ltd (United Kingdom) based on the colorimetric method described by [20], where creatinine reacts with picrate in an alkaline solution to form a colored complex. The reagents used included a standard at 177 μ mol/L, picric acid at 35 mmol/L, sodium hydroxide at 1.6 mmol/L, and trichloroacetic acid (TCA) at 1.2 mol/L. For the assay, 1.0 mL of TCA and 1.0 mL of serum were pipetted into centrifuge tubes, mixed with a glass rod to evenly disperse the precipitate, and centrifuged at 2,500 rpm for 10 minutes. The supernatant was collected for further analysis. In the assay procedure, 0.5 mL of distilled water was used for the blank, 0.5 mL of TCA for the standard, and 1.0 mL of supernatant for the sample, all mixed with 1.0 mL of reagent mixture. Each tube was incubated for 20 minutes at 37°C, and the absorbance of the sample (A_sample) and standard (A_standard) was measured against the blank at 520 nm.

Concentration of creatinine in serum is given as =

$$\frac{A \text{ sample} * 177}{A \text{ standard}} \mu\text{mol/L}$$

2.6.12 Blood urea nitrogen (BUN)

Blood urea nitrogen was analyzed using the standard procedure from Randox Laboratory Ltd (United Kingdom), based on the colorimetric method outlined by [21]. Urea in the serum is hydrolyzed to ammonia in the presence of urease, and the ammonia is then measured photometrically by Berthelot's reaction. Sodium nitroprusside and urease were combined as solution 2, phenol was diluted with 660 mL of distilled water as solution 3, and sodium hypochlorite was diluted with 750 mL of distilled water as solution 4. 10 μ L of distilled water was used for the blank, 10 μ L of the standard solution for the standard, and 10 μ L of the sample for the sample, all mixed with 100 μ L of solution 2 [22-25]. This mixture was incubated at 37°C for 10 minutes. Subsequently, 2.50 mL of solution 3 and 2.50 mL of solution 4 were added to each tube, mixed, and incubated again at 37°C for 15 minutes. The absorbance of the sample (A sample) and standard (A standard) was measured against the blank at 546 nm. The urea concentration was calculated using the following method [26-29].

$$\text{Urea Concentration} = \frac{A_{\text{sample}} \times 80}{A_{\text{standard}}} \text{ (mg/dl)}$$

Where 1 mg of urea corresponds to 0.467mg of urea nitrogen [30-32].

3. RESULTS AND DISCUSSION

3.1 Physical Observations

The following were observed during the course of this project

- I. Reduced weight gain, excessive thirst, hypo activity, lack of coordination, lethargy, difficulty breathing, emaciation, and prostration in groups C and D. Three rats died in group C and two in group D.
- II. During the process of sacrifice, fat deposits were seen around the kidneys in the rats given aqueous ginger extract, with more deposits observed in group B compared to group D [33,34].

3.2 Histopathological Evaluation

H/E = Hematoxylin and eosin staining

Kidney GA (X400 Magnification): The H&E-stained kidney sections at 400x magnification showed normal renal glomeruli and tubules. However, there was evidence of congestion in the cortical vessels and tubules, indicating possible circulatory disturbances or inflammatory processes [34].

Kidney GB (X400 Magnification): H/E staining shows normal renal glomeruli and tubules. There is moderate congestion of the medullary vessels.

Kidney GC (X400 Magnification): H/E staining shows preserved renal tissue, with widened Bowman's spaces and collecting ducts. The collecting ducts are frequently distorted, seen in both longitudinal and transverse sections. Some of the ducts contain eosinophilic secretions.

Kidney Gd (X400 Magnification): H/E staining shows severe architectural distortion. There is diffuse cortical and medullary congestion extending to involve the glomeruli. There is also severe interstitial nephritis.




Fig. 2. Plates of histopathological evaluation

3.3 Summary of Histopathological Analysis

Table 1. Summary of hispathological analysis of the rat kidneys

Groups	Renal Architecture	Cortex	Medulla	Glomerulli	Tubules	Interstitium
Group A	Preserved	Congested		-	-	-
Group B	Preserved	-	Congested	-	-	-
Group C	ivliid distortion			WideBM spaces	Distorted	Severe nephritis
Group D	Severe distortion	Congested	Congested	Congested	Distorted	Severe nephritis

Table 2. Mortality effect of treatment on rats

Groups	No of Rats	Dosage of Gentamacin administration	Dosage of Ginger Administration	No of Death
Group A	6	100mg/kg	-	-
Group B	6	100mg/kg	2.14ml/kg	-
Group C	9		-	3
Group D	8		2.14ml/kg	2

3.4 Biochemical Evaluation

Table 3. Effect of treatment on PCV and WBC count

Female Albino Rats	Mortality (N)	Packed cell Volume (PCV)%	White blood cell (WBC/L)x10 ⁶
Group A	NIL	41.8	1.1
Group B	NIL	44.4 ^a	1.03 ^a
Group C	3	35 ^{ab}	0.96 ^{a,b}
Group D	2	33.5 ^{abx}	0.7.8 ^{a,b,c}

Table 4. Effect of treatment on serum albumin creatinine and bun

Female Albino Rats	Serum Creatinine (mg/L)	Serum Albumin (g/L)	Blood Urea Nitrogen level (BUN). (g/l)
Group A	0.74	51.2	17.82
Group B	1.11 ^a	37.29a	12.95 ^a
Group C	1.6 ^{a,b}	35.96 ^{a,n}	122.52 ^{a,b}
Group D	1.9 ^{a,b,c}	35.65 ^{a,n,x}	45.33 ^{ab,c}

Table 5. Effect of treatment on total cholesterol and triglyceride

Female Albino Rats	Serum total cholesterol mg/dL	Serum Triglyceride
Group A	189.1	163.4
Group B	168.4 ^a	108.6 ^a
Group C	69.8 ^{a,b}	182.8 ^{n,b}
Group D	39.5 ^{a,b,c}	79.9 ^{a,nb,c}

Table 6. Effect of treatment on some antioxidant enzymes and oxidative stress parameters

Female Albino Rats	GSH Level (U/L) x 10 ⁻⁴	GST Activity (U/L) X 10 ⁻³	SOD Activity (U/L) x 10 ⁻⁴	Catalase Activity (U/L) x 10 ⁻²	MDA Level (n mol/MI) x 10 ⁻⁵
Group A	3.31	11.7	2.2	.732	1.7
Group B	3.0 ^{na}	5.0 ^a	2.4 ^{na}	4.3 ^a	1.98 ⁿ
Group C	2.2 ^{a,b}	5.21 ^{a,nb}	0 ^{a,b}	5.5 ^{a,n}	2.4 ^{a,b}
Group D	2.9 ^{a,nb,c}	5.5 ^{a,nb,nc}	0 ^{a,b,nc}	3.4 ^{a,n,x}	2.5 ^{a,b,nc}

3.5 Graphical Representation of Biochemical Evaluation




Fig. 3. Graph Showing Effect of Treatment of Packed Cell Volume (PCV) of Rats Blood




Fig. 4. Graph showing effect of treatment on white blood cell (WBC) of rats blood ($\times 10^9$ per Litre)




Fig. 5. Graph showing effect of treatment on creatinine level of blood serum of rat (Mg/L)




Fig. 6. Graph showing effect of treatment on serum albumin of rat (g/L)




Fig. 7. Graph showing effect of treatment on the blood urine nitrogen (BUN) level in the serum of rate (g/L)




Fig. 8. Graph showing effect of treatment on total cholesterol rat serum (mg/L)




Fig. 9. Graph showing effect of treatment on triglycerides of rat blood serum (mg/dl)




Fig. 10. Graph showing effect of treatment on reduction glutathione (GSH) in rat kidney (Homogenate) (μ /L)




Fig. 11. Graph showing effect of treatment on glutathione-s-transferase (GST) in rat kidney (Homogenate) (μL)




Fig. 12. Graph showing effect of treatment on the activity of superoxide dismutase (SOD) in rat kidney homogenate (μL)




Fig. 13. Graph showing effect of treatment on the catalase activity in rat kidney (Homogenate) (μL)




Fig. 14. Graph showing effect of treatment on malondialdehyde (MDA) (Nmol/ml)

3.6 Summary of Results

Packed Cell Volume (PCV): From Table 3, there is a significant difference in PCV between groups B, C, D and the normal control group A. PCV is significantly lower ($p < 0.05$) in group A compared to group B, and significantly higher compared to groups C and D. PCV is significantly higher in group B compared to groups C and D, and significantly higher in group C compared to group B.

White Blood Cell Count (WBC): As shown in Table 3, the white blood cell levels were observed to significantly ($p < 0.05$) decrease in groups B, C, and D compared to group A. WBC is significantly higher in group B compared to groups C and D, and significantly higher in group C compared to group D.

Serum creatinine: The effect of treatment on creatinine level was observed in all the groups, as shown in Table 4. The creatinine level significantly increased in groups B, C, and D compared to group A. The creatinine levels in groups C and D significantly increased compared to group B. The creatinine level in group A also significantly increased compared to group C.

Serum albumin: The effect of treatment on serum albumin was observed in all the groups. There was a significant decrease in serum albumin in groups B, C, and D compared to group A. There was no significant difference in serum albumin between group C and group B, nor was there any significant difference between group D and group C.

Blood urea nitrogen (BUN): The level of blood urea nitrogen (BUN) was measured in the different groups, as summarized in Table 4. The BUN level was significantly ($p < 0.05$) higher in group A compared to group B, and lower in group A compared to groups C and D. The BUN level significantly ($p < 0.05$) decreased in group D compared to group C.

Total serum cholesterol: The level of total serum cholesterol was measured in the different groups, as shown in Table 5. The cholesterol level in group A was significantly ($p < 0.05$) higher compared to groups B, C, and D. The cholesterol level in group B was significantly ($p < 0.05$) higher compared to groups C and D, while group C was significantly ($p < 0.05$) higher compared to group D.

Triglycerides: The level of triglycerides in all groups was also measured, as shown in Table 5. There was a significant decrease in the triglyceride level in groups B (108.8) and D (79.9) compared to group A (163.4), while group C (182.8) showed a significant ($p < 0.05$) increase compared to group A. Group B showed a significant ($p < 0.05$) decrease compared to group C, with no significant difference compared to group D. Group D showed a significant decrease compared to group C.

Catalase activity: The catalase activity was measured in all groups, as shown in Table 6. The level of catalase activity significantly ($p < 0.05$) decreased in group A (0.007) compared to groups B (0.044), C (0.055), and D (0.034). There was no significant ($p > 0.05$) difference in catalase activity between group B and groups C and D, while group D showed a significant ($p > 0.05$) decrease in catalase activity compared to group C.

Glutathione Level (GSH): The level of GSH was measured in all the groups, as shown in Table 6. The observations showed no significant difference in the GSH level between group A and group B. There was a significant decrease in GSH level in groups C and D compared to group A. The GSH level significantly decreased in groups C and D compared to group B. The GSH level significantly increased in group B compared to group C.

Glutathione-S-Transferase (GST): The activity of GST was measured in all the groups, as shown in Table 6. The observation shows a significant ($p < 0.05$) decrease in groups B, C, and D compared to group A. There was no significant difference in GST activity between group B and groups C and B, nor was there any significant difference between group C and group D.

Malondialdehyde Assay (MDA): The level of malondialdehyde was measured in all the groups, as shown in Table 6. The MDA level increased in groups B, C, and D. There was no significant increase in MDA in group B compared to group A, while the MDA level increased in groups C and D compared to group B. When compared to group B, there was a significant difference in the MDA level in groups C and D. However, there was no significant difference between group C and group D.

Superoxide Dismutase (SOD): The activity of SOD was measured, as shown in Table 6. The statistical analysis shows no significant ($p > 0.05$) difference between group A (normal control) and group B. The SOD activity significantly decreased in groups C and D compared to group A. It also shows a significant decrease in SOD activity in groups C and D compared- to group B. SOD activity significantly ($p < 0.05$) decreased in group C compared to group B.

3.7 Discussion

Gentamicin, an aminoglycoside antibiotic widely used for treating serious bacterial infections, has been associated with acute kidney injury as a significant adverse effect. Numerous studies have established that reactive oxygen species (ROS) play a pivotal role in the nephrotoxicity induced by gentamicin. The accumulation of ROS leads to oxidative stress, which is a key contributor to cellular damage in renal tissues [35]. Administration of gentamicin has been shown to result in substantial nephrotoxicity, characterized by increased levels of blood urea nitrogen (BUN) and creatinine, alongside a marked reduction in glutathione (GSH) levels and an increase in malondialdehyde (MDA) [36]. The elevation of BUN and creatinine levels indicates impaired renal function, while the decrease in GSH and increase in MDA reflect oxidative damage to kidney cells [19]. The tissue damage induced by gentamicin is primarily mediated through oxidative mechanisms, highlighting the importance of ROS in the pathophysiology of gentamicin-induced nephrotoxicity [37,38]. Ginger has demonstrated strong antioxidant properties, with its component gingerone inhibiting lipid peroxidation and effectively scavenging free radicals [39,40]. Animal studies have suggested that ginger can reduce cholesterol levels and positively impact plasma lipid composition, potentially preventing atherosclerosis [39,40]. However, the effects of ginger on gentamicin-induced nephrotoxicity are mixed. While ginger has antioxidant properties, some studies found that concurrent administration of ginger and gentamicin may have cumulative negative effects, reducing immunity and exacerbating kidney damage [38]. Ginger administration, either separately or simultaneously with gentamicin, significantly increased MDA levels, suggesting that ginger extract may worsen gentamicin-induced nephrotoxicity [39]. Histopathological analysis showed more severe architectural distortion and

interstitial nephritis in rats receiving both gentamicin and ginger compared to gentamicin alone [37,39]. Creatinine levels, a major marker of kidney dysfunction, were significantly increased in groups receiving ginger, indicating that ginger may enhance gentamicin-induced nephrotoxicity rather than mitigate it [37]. While ginger has antioxidant properties and potential benefits for reducing cholesterol and preventing atherosclerosis, its effects on gentamicin-induced nephrotoxicity are unclear. Some studies suggest ginger may exacerbate kidney damage caused by gentamicin, while others indicate potential protective effects. Further research is needed to clarify the relationship between ginger and gentamicin-induced nephrotoxicity.

4. CONCLUSION

In conclusion, this research indicates that the aqueous extract of ginger did not provide a protective benefit against oxidative stress and nephrotoxicity induced by gentamicin. This lack of protection may be attributed to factors such as dosage levels or the catalytic activity of iron content in ginger, which could potentially exacerbate oxidative damage rather than mitigate it. Despite these findings, it is important to highlight the significant health benefits associated with ginger extract, particularly its ability to lower serum cholesterol levels as elevated cholesterol is a well-known risk factor for atherosclerotic conditions and cardiovascular diseases. Also ginger's rich composition of bioactive compounds, including gingerols and shogaols, contributes to its well-documented antioxidant, anti-inflammatory, and anti-hyperlipidemic properties. These attributes suggest that ginger could play a multifaceted role in health promotion beyond its use as a culinary spice. Future studies should explore optimal extraction methods, dosages, and formulations to fully harness the therapeutic potential of ginger.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ETHICAL APPROVAL

Animal ethic committee approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Martin F. Fundamentals of Anatomy and Physiology 5th Edition. Prentice hall international Inc; 2001.
2. Coe FL, Eva A, Buorchester E. Kidney Stone disease. J. Clin invest. 2005; 115(10):2598-608.
3. Goldfarb BS, Coe FL. Prevention of recurrent nephrolithiasis Amfam physician. 1999;60(8):2269-76.
4. Needleman H. Lead poisoning. Annu Rev Med. 2004;55:209-22.
5. Prakash J, Shrin P, Adel R. Chemical composition and antioxidant properties of ginger root. J Med PI Res. 2010;4(24): 2674-2679.
6. Zancan KC, Marques MO, Petenate AJ, Meireles MA. Extraction of ginger (*Zingiber officinale* Roscoe) oleoresin with CO₂ and co-solvents: A study of the antioxidant action of the extracts. J Supercrit Flu. 2002;24:57-76.
7. Maghsoudi S, Gol A, Dabiri S, Javadi A. Preventive effect of ginger (*Zingiber officinale*) pretreatment on renal ischemia-reperfusion in rats. Eur Surg Res. 2011;46:45–51.
8. Wilkinson JM. Effect of ginger tea on fatal development of sprague dawley rats reprod toxicol. 2000;14(6):507-512.
9. Gharahi GH, Shafiee M, Zare M. The protective effect of ginger against malathion-induced oxidative stress in rat liver. Environmental Toxicology and Pharmacology. 2016;43:95-100.
10. Bae JH, Kim HJ, Lee SK. Antitumor effects of ginger and its bioactive compounds in cancer cells: A review. Phytotherapy Research. 2019;33(10):2671-2681.
11. Zhang Q, Li Y. The anti-inflammatory effects of 6-gingerol in mouse models. Journal of Agricultural and Food Chemistry. 2017;65(12):2531-2537.
12. Alam MM, Ullah MF, Khan MI. Neuroprotective effects of ginger (*Zingiber officinale*) and its bioactive components in neurodegenerative diseases: A review. Journal of Ethnopharmacology. 2020;258: 112867.
13. Khan MA, Khedher NB. Anti-inflammatory effects of ginger oil in a rat model of arthritis. Journal of Medicinal Food. 2018; 21(4):391-397.
14. Garza-Cadena C, Ortega-Rivera DM, Machorro-García G, Gonzalez-Zermeño EM, Homma-Dueñas D, Plata-Gryl M, Castro-Muñoz R. A comprehensive review on ginger (*Zingiber officinale*) as a potential source of nutraceuticals for food formulations: Towards the polishing of gingerol and other present biomolecules. Food Chemistry. 2023;413:135629.
15. Maghraby YR, Labib RM, Sobeh M, Farag MA. Gingerols and shogaols: A multi-faceted review of their extraction, formulation, and analysis in drugs and biofluids to maximize their nutraceutical and pharmaceutical applications; 2023.
16. Shaukat MN, Nazir A, Fallico B. Ginger bioactives: A comprehensive review of health benefits and potential food applications; 2023.
17. Cruz MA, Oliveira AC, Silva LM. Anesthesia and analgesia in laboratory animals: A review. Journal of the American Association for Laboratory Animal Science. 2020;59(5):555-564.
18. Hoffman RM, Ritchie RF, Houghton JD. Cardiac puncture: A technique for blood collection in small laboratory animals. Journal of Laboratory Animal Science. 2018;8(4):245-250.
19. Friedman AD, Reddy SM, Weller DR. Optimizing serum collection and processing for clinical chemistry analysis. Clinical Biochemistry. 2021;84: 1-8.
20. Kumar A, Gupta R, Sharma S. Biochemical markers in clinical practice: A review of their significance in health and disease. Journal of Clinical Laboratory Analysis. 2021;35(3):e23725.
21. Baker FJ, Silverton. Introduction to Medical Laboratory Technology. Fifth edition. 1984;556-579.
22. Hildebraunt AG, Roots I. Reduced nicotinamide adenine dinucleotide phosphate (NADPH)-dependent formation and breakdown of hydrogen peroxide during mixed function oxidation reactions in liver microsomes. Arch Biochem Biophys. 1975, Dec;171(2):385-97.
23. Marlund S, Marklund G. Involvement of superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. Eur. J. Biochem. 1974;1974:47 469-74.

24. Buege JA, Aust SD. Microsomal lipid peroxidation. *Methods in Enzymology*. 1978;52:302-310.
25. Wills ED. *Biochem J*. 1969;113:315.
26. Habig WH, Pabst MJ, Jakoby WB. Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *J Biol Chem*. 1974, Nov 25;249(22):7130-9. PMID: 4436300.
27. Carlberg I, Mannervik B. Glutathione reductase. *Methods Enzymol*. 1985;113:484-90.
DOI: 10.1016/s0076-6879(85)13062-4.
PMID: 3003504.
28. Boyland E. Reduced glutathione assay. In *Methods in Enzymology*. Academic Press. 2001;348:346-348.
29. Trinder P. Enzymatic determination of glucose in blood serum. *Annals of Clinical Biochemistry*. 1969;6:24.
30. Richmond W. *Clinician Chem*. 1973;19:1350.
31. Rodkey FL. *Clin., Chem*. 1964;10:606.
32. Henry RJ. *Clinical Chemistry, principles and Technics*, 2nd Edition. 1974;525.
33. Weatherbum MW. *Anal Chem*. 1967;39:971.
34. Junqueira, LC, Carneiro J. *Basic histology: Text and atlas* (13th ed.). McGraw-Hill Education; 2013.
35. Al-kuraishy HM et al. Gentamicin-induced nephrotoxicity: Mechanisms and protective strategies. *Journal of Nephrology*. 2019;32(6):919-930.
36. Balakumar P et al. Gentamicin-induced nephrotoxicity: Mechanisms and protective strategies. *Pharmacological Reports*. 2010;62(4):667-677.
37. Sener Goskel, Sehir u, A. Ozer et al. *Journal of Pineal Research*. 2002, May;32(4):231-236(6).
38. Krishnakantha T, Lokesh B. scavenging of superoxide anions by spice principle. *Indian J. biochem Biophys*. 1993;30(2):133-134.
39. Antoine Al-Achi. *A Current Look at Ginger use*, Campbell University, School of Pharmacy, Buies creek, NC; 2007.
40. Rosert J. Drug-induced acute interstitial nephritis. *Kidney Int*. 2001;60(2):804-17.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/121844>