



A Promising Sustainable Pest Management Technology: Microbial Bio-Control Agent

**Utpal Dey ^{a*}, Shatabhisa Sarkar ^a, Meenakshi Malik ^b, Mukesh Sehgal ^b,
Pijush Debbarma ^a and Subhash Chander ^b**

^a *Krishi Vigyan Kendra, Sepahijala, CAU (I), Tripura, India.*

^b *ICAR-National Research Centre for Integrated Pest Management, New Delhi, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/MRJI/2021/v31i1230361

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/69080>

Short Communication

Received 22 April 2021
Accepted 29 June 2021
Published 30 December 2021

ABSTRACT

Microbial bio-control agents play an important role in integrated pest management in modern agriculture for managing pests without affecting environment and humans. Recently, microbial bio-control agents are gaining more attention as they are easy to use, safe and an alternative eco-friendly approach of hazardous pesticide chemicals. Biological control agents are ecologically sustainable and effective crop protection approach in agriculture and horticultural crops including organic systems. Predator, parasitism or other natural mechanism can be use to reduce the pest such as insects, mites, weeds and plant diseases.

Keywords: Crop protection, microbial bio-control agents, natural mechanism, parasitism.

1. INTRODUCTION

Biological control agents are ecologically sustainable and effective crop protection approach in agriculture and horticultural crops including organic systems. It is one of the most

promising components of an integrated pest management strategies [1,2]. Generally, biological agent refers the use of living organism or living things to reduce key pest. Biological agent feed on the pest and thus suppresses the pest population [3].

*Corresponding author: E-mail: utpaldey86@gmail.com;

The biological process in environment is rarely very complex as the relation / interaction between plants, animal and other living or non-living organism in the environment that make balance in the system. Natural process that they depend on each other feeding on lower organism to higher organism and decomposer etc. and vice-versa. Predator, parasitism or other natural mechanism can be used to reduce key pest such as insects, mites, weeds and plant diseases. In this case involve an active human management.

It is an economical, sustainable, reliable and environmentally friendly pest management method by using living organisms to suppress the numbers of key pest species. The adverse effect on the environment, human being, livestock etc is minimized by reducing the dependence on chemical pesticides [4]. It increases species diversity and conserves it within an agro-ecosystem. Biological control can also be considered as a potential tool for achieving sustainable integrated pest management in organic/natural farming system. Microbial bio-control agents are applied in agricultural and horticultural crops to suppress the plant pathogens [5].

1.1 Types of Microbial Biocontrol Agents

Natural antagonistic microorganisms are responsible to reduces plant pests. Different types of microbes viz., bacteria, fungi, viruses, and protozoans etc has enormous potential as biocontrol agents and alternative approach of chemical-based pesticides. Some of the fungal biocontrol agents are *Trichoderma harzianum*, *Trichoderma viride*, *Verticillium lecanii*, *Metarhizium anisopliae*, *Paecilomyces lilacinus* etc. Bacterial biocontrol agents viz., *Pseudomonas fluorescens*, *Bacillus subtilis*, *Bacillus thuringiensis* also plays an important role in suppression of soil-borne plant pathogenic organisms.

1.2 Mode of action

Antibiosis, Mycoparasitism / Hyperparasitism, Competition, Lytic enzymes, Hydrogen cyanide, Induced Systemic Resistance and Plant growth promotion etc [6]. This process is quite slow but effective, environmentally suitable, does not pollute and toxic to other organism than specific.

2. MODE OF APPLICATION

2.1 Seed Treatment

Seeds can be treated with *Trichoderma harzianum* or *T. viride* @ 5-10 g/kg seed or *Pseudomonas fluorescens* @ 5-10 g/kg seed or *Beauveria bassiana* or *Metarhizium anisopliae* @ 10 g/kg seed or *Verticillium lecanii* @ 20 ml/kg or *Paecilomyces lilacinus* @ 4-6 g/kg seed or *Bacillus subtilis* @ 10 g/kg seed and kept for shade drying for 30 minutes.

2.2 Nursery Treatment

Drench nursery bed with *Trichoderma* @ 5g/L or 10-25g of *Trichoderma* per 100 square meter area or *Pseudomonas fluorescens* 50 -100g formulation per square meter area or *Paecilomyces lilacinus* 50g formulation per square meter area.

2.3 Seedling Treatment

Roots of the seedlings are dipped into *Trichoderma* solution (20 – 25 g/L) or *Pseudomonas fluorescens* solution @ 5g/L or *Verticillium lecanii* solution @ 20 ml/L or *Paecilomyces lilacinus* solution @ 10 g/L or *Bacillus subtilis* solution @ 5g/L for 20-30 minutes before transplanting.

2.4 Soil Application

Apply 4-5 kg *Trichoderma* or 2.5 kg *Pseudomonas fluorescens* or *B. bassiana* or *M. anisopliae* @ 10 kg/ha or *Verticillium lecanii* @ 1L/acre or *Paecilomyces lilacinus* @ 2kg/acre or *Bacillus thuringiensis* @ 1.12kg/ha or 2.5-5 kg *Bacillus subtilis* with 100kg decomposed manure/farm yard manure 20-25 days before applying in field and cover it with polythene sheet. Sprinkle the heap with little water immediately. Turn the mixture in every 3-4 days interval.

2.5 Foliar Application

5 kg/ hectare in 500 liters of water i.e., 5 gm per liter of water.

2.6 Advantages

Biological control agents are eco-friendly approach and do not have any adverse effect on the environment, human being, livestock etc [7]. Comparatively cheaper than other agrochemical like pesticides and insecticides. These methods



Fig. 1. Demonstration of biocontrol agents at Pathalia village, Sepahijala Tripura



Fig. 2. Paddy seedlings root dip method



Fig. 3. Treated seedlings transplanted in lines



Fig. 4. Method demonstration on biocontrol agents in farmers field



Fig. 5. Method demonstration on biocontrol agents in farmers field



Fig. 6. Seed treatment with biocontrol agents

are also easy to use, attack only target organism, readily available and can be used in any season. The most crucial use of biocontrol agents gives protection to crops throughout the cropping season

and help in reducing the use of chemicals and pesticides which have a harmful effect on human beings [8].

3. CONCLUSION

Microbial biocontrol agents play an important role in integrated pest management in modern agriculture reducing pests without affecting environment and human. Recently, microbial biocontrol agents gaining more attention as they are easy to use, safe and an alternative eco-friendly approach of chemicals.

4. LIMITATIONS

High specificity against the target pest resulting multiple microbial pesticides to be used. The biocontrol agents do not eradicate all the pests and are a useful and economical tool for pest control only when used on a large scale.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Gurr GM, Wratten SD, Landis DA, You M. Habitat management to suppress pest populations: Progress and prospects. *Annual Review of Entomology*. 2017; 62:91–109.
2. Michaud J. Challenges to conservation biological control on the High Plains: 150 years of evolutionary rescue. *Biological Control*. 2018; 125:65– 73.
3. Heimpel GE, Mills N. *Biological Control - Ecology and Applications*. Cambridge: Cambridge University Press. 2017.
4. Damalas Christos A, Eleftherohorinos Ilias G. Pesticide Exposure, Safety Issues, and Risk Assessment Indicators. *Int. J. Environ. Res. Public Health*. 2011;8(5): 1402-1419.
5. Singh S, Kumar V, Datta S, Singh S, Dhanjal DS, Dhaka V, Sharma K, Singh J. Commercial production and formulation of microbial biocontrol agents. *Food Security and Plant Disease Management*; 2021.
6. Köhl J, Kolnaar R, Ravensberg WJ. Mode of action of microbial biological control agents against plant diseases: Relevance beyond efficacy. *Frontiers in Plant Science*. 2019; 10:845.
7. Syed-Ab-Rahman, Sharifah Farhana, Singh Eugenie, Pieterse Corné, Schenk Peer. Emerging Microbial Biocontrol Strategies for Plant Pathogens. *Plant Science*. 2017; 267.
8. Peshin R, Bandral RS, Zhang W, Wilson L, Dhawan AK. Integrated Pest Management: A Global Overview of History, Programs and Adoption. In *Integrated Pest Management: Innovation-Development Process*; Peshin, R., Dhawan, A.K., Eds.; Springer: Dordrecht, The Netherland, 2009; 1–50.

© 2021 Dey et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/69080>