

Research Article

Computations of the M-Polynomials and Degree-Based Topological Indices for Dendrimers and Polyomino Chains

Young Chel Kwun ¹, Adeel Farooq,² Waqas Nazeer ³, Zohaib Zahid,⁴
Saba Noreen,⁵ and Shin Min Kang ^{6,7}

¹Department of Mathematics, Dong-A University, Busan 49315, Republic of Korea

²Department of Mathematics, COMSATS University Islamabad, Lahore Campus, Lahore 54000, Pakistan

³Division of Science and Technology, University of Education, Lahore 54000, Pakistan

⁴Department of Mathematics, University of Management and Technology, Lahore 54000, Pakistan

⁵Department of Mathematics and Statistics, The University of Lahore, Lahore, Pakistan

⁶Department of Mathematics and RINS, Gyeongsang National University, Jinju 52828, Republic of Korea

⁷Center for General Education, China Medical University, Taichung 40402, Taiwan

Correspondence should be addressed to Waqas Nazeer; nazeer.waqas@ue.edu.pk and Shin Min Kang; smkang@gnu.ac.kr

Received 16 July 2018; Accepted 18 October 2018; Published 1 November 2018

Academic Editor: Charles L. Wilkins

Copyright © 2018 Young Chel Kwun et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Topological indices correlate certain physicochemical properties like boiling point, stability, and strain energy of chemical compounds. In this report, we compute M-polynomials for PAMAM dendrimers and polyomino chains. Moreover, by applying calculus, we compute nine important topological indices of under-study dendrimers and chains.

1. Introduction

The **polyomino chains** constitute a finite 2-connected floor plan, where each inner face (or a unit) is surrounded by a square of length one. We can say that it is a union of cells connected by edges in a planar square lattice. For the origin of dominoes, we quote [1]. The **polyomino chains** have a long history dating back to the beginning of the 20th century, but they were originally promoted by Golomb [2, 3]. Dendrimers [4] are repetitively branched molecules. The name comes from the Greek word, which translates to “trees.” Synonymous terms for dendrimers include arborols and cascade molecules. The first dendrimer was made by Fritz Vögtle in [5]. For detailed study about dendrimer structures we refer the reader to [6–9].

Many studies have shown that there is a strong intrinsic link between the chemical properties of chemical compounds and drugs (such as melting point and boiling point) and their molecular structure [10, 11]. The topological index

defined on the structure of these chemical molecules can help researchers better understand the physical characteristics, chemical reactivity, and biological activity [12]. Therefore, the study of topological indices of chemical substances and chemical structures of drugs can make up for the lack of chemical experiments and provide theoretical basis for the preparation of drugs and chemical substances. In the previous two decades, a number of topological indices have been characterized and utilized for correlation analysis in pharmacology, environmental chemistry, toxicology, and theoretical chemistry [13]. Hosoya polynomial (Wiener polynomial) [14] plays a pivotal role in finding topological indices that depend on distances. From this polynomial, a long list of distance-based topological indices can be easily evaluated. A similar breakthrough was obtained recently by Klavžar et al. [15], in the context of degree-based indices. In the year 2015, authors in [15] introduced the M-polynomial, which plays similar role “to what Hosoya polynomial does” to determine many topological indices depending on the degree of end vertices [16–20].

In the present paper, we compute M-polynomials for different dendrimer structures and polyomino chains. By applying fundamental calculus, we recover nine degree-based topological indices for these dendrimers and chains.

2. Basic Definitions and Literature Review

In this paper, we fixed G as a connected graph, $V(G)$ is the set of vertices, $E(G)$ is the set of edges, and d_v is the degree of any vertex v . Most of the definitions presented in this section can be found in [17].

Definition 1 (see [15]). The M-polynomial of G is defined as

$$M(G; x, y) = \sum_{\delta \leq i \leq j \leq \Delta} m_{ij}(G) x^i y^j \quad (1)$$

where $\delta = \min\{d_v \mid v \in V(G)\}$, $\Delta = \max\{d_v \mid v \in V(G)\}$, and $m_{ij}(G)$ is the edge $vu \in E(G)$ that is $i \leq j$.

The very first topological index was the Wiener index, defined by Wiener in 1945, when he was studying boiling point of alkane [21]. For comprehensive details about the applications of Wiener index, see [22, 23]. After that, in 1975, Milan Randić [24] introduced the first degree-based topological index, which is now known as Randić index and is defined as

$$R_{-1/2}(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_u d_v}}. \quad (2)$$

The generalized Randić index is defined as

$$R_\alpha(G) = \sum_{uv \in E(G)} \frac{1}{(d_u d_v)^\alpha}; \quad (3)$$

please see [25–29].

The inverse generalized Randić index is defined as

$$RR_\alpha(G) = \sum_{uv \in E(G)} (d_u d_v)^\alpha. \quad (4)$$

It can be seen easily that the Randić index is particular case of the generalized Randić index and the inverse generalized Randić index. Other oldest degree-based topological indices are Zagreb indices. The first Zagreb index is defined as

$$M_1(G) = \sum_{uv \in E(G)} (d_u + d_v) \quad (5)$$

and the second Zagreb index is defined as

$$M_2(G) = \sum_{uv \in E(G)} (d_u \times d_v). \quad (6)$$

The second modified Zagreb index is defined as

$${}^m M_2(G) = \sum_{uv \in E(G)} \frac{1}{d(u) d(v)}. \quad (7)$$

For detailed study about Zagreb indices, we refer the reader to [30–32]. There are many other degree-based topological indices, for example, symmetric division index:

$$SDD(G) = \sum_{uv \in E(G)} \left\{ \frac{\min(d_u, d_v)}{\max(d_u, d_v)} + \frac{\max(d_u, d_v)}{\min(d_u, d_v)} \right\} \quad (8)$$

harmonic index:

$$H(G) = \sum_{vu \in E(G)} \frac{2}{d_u + d_v} \quad (9)$$

inverse sum index:

$$I(G) = \sum_{vu \in E(G)} \frac{d_u d_v}{d_u + d_v} \quad (10)$$

augmented Zagreb index:

$$A(G) = \sum_{vu \in E(G)} \left\{ \frac{d_u d_v}{d_u + d_v - 2} \right\}^3. \quad (11)$$

We refer to [33–45] for detailed survey about the above defined indices and applications. Tables exhibited in [15–19] relate some notable degree-based topological indices with M-polynomial with the following notations [17]:

$$\begin{aligned} D_x &= x \frac{\partial(f(x, y))}{\partial x}, \\ D_y &= y \frac{\partial(f(x, y))}{\partial y}, \\ S_x &= \int_0^x \frac{f(t, y)}{t} dt, \\ S_y &= \int_0^y \frac{f(x, t)}{t} dt, \\ J(f(x, y)) &= f(x, x), \\ Q_\alpha(f(x, y)) &= x^\alpha f(x, y). \end{aligned} \quad (12)$$

3. Computational Results

In this section we give our computational results.

3.1. M-Polynomials and Degree-Based Indices for PAMAM Dendrimers. Polyamidoamine (PAMAM) dendrimers are hyperbranched polymers with unparalleled molecular uniformity, narrow molecular weight distribution, defined size and shape characteristics, and a multifunctional terminal surface. These nanoscale polymers consist of an ethylenediamine core, a repetitive branching amidoamine internal structure, and a primary amine terminal surface. Dendrimers are “grown” off a central core in an iterative manufacturing process, with each subsequent step representing a new “generation” of dendrimer. Increasing generations (molecular weight) produce larger molecular diameters, twice the

number of reactive surface sites and approximately double the molecular weight of the preceding generation. PAMAM dendrimers also assume a spheroidal, globular shape at generation 4 and above (see molecular simulation below). Their functionality is readily tailored, and their uniformity, size, and highly reactive “molecular Velcro” surfaces are the functional keys to their use. Here we consider PD_1 , which denote PAMAM dendrimers with trifunctional core unit generated by dendrimer generations G_n with n growth stages, and PD_2 , the PAMAM dendrimers with different core generated by dendrimer generators G_n with n growth stages. DS_1 is kinds of PAMAM dendrimers with n growth stages

Theorem 2. For the PAMAM dendrimers PD_1 , we have

$$\begin{aligned} M(PD_1, x, y) &= 3 \cdot 2^n xy^2 + 3(2^{n+1} - 1)xy^3 \\ &\quad + 9(2^{n+1} - 1)x^2y^2 \\ &\quad + 3(7 \cdot 2^n - 4)x^2y^3. \end{aligned} \quad (13)$$

Proof. Let PD_1 denote PAMAM dendrimers with trifunctional core unit generated by dendrimer generations G_n with n growth stages.

The edge set of PD_1 has following four partitions:

$$\begin{aligned} E_{\{1,2\}} &= \{e = uv \in E(PD_1) \mid d_u = 1, d_v = 2\}, \\ E_{\{1,3\}} &= \{e = uv \in E(PD_1) \mid d_u = 1, d_v = 3\}, \\ E_{\{2,2\}} &= \{e = uv \in E(PD_1) \mid d_u = 2, d_v = 2\}, \\ E_{\{2,3\}} &= \{e = uv \in E(PD_1) \mid d_u = 2, d_v = 3\}. \end{aligned} \quad (14)$$

Now

$$\begin{aligned} |E_{\{1,2\}}| &= 3 \cdot 2^n, \\ |E_{\{1,3\}}| &= 6 \cdot 2^n - 3, \\ |E_{\{2,2\}}| &= 18 \cdot 2^n - 9, \end{aligned} \quad (15)$$

and

$$|E_{\{2,3\}}| = 21 \cdot 2^n - 12.$$

$$\begin{aligned} M(PD_1; x, y) &= \sum_{i \leq j} m_{ij}(PD_1) x^i y^j \\ &= \sum_{1 \leq 2} m_{12}(PD_1) xy^2 \\ &\quad + \sum_{1 \leq 3} m_{13}(PD_1) x^1 y^3 \\ &\quad + \sum_{2 \leq 2} m_{22}(PD_1) x^2 y^2 \\ &\quad + \sum_{2 \leq 3} m_{23}(PD_1) x^2 y^3 \\ &= \sum_{uv \in E_{\{1,2\}}} m_{12}(PD_1) xy^2 \\ &\quad + \sum_{uv \in E_{\{1,3\}}} m_{13}(PD_1) xy^3 \end{aligned}$$

$$\begin{aligned} &\quad + \sum_{uv \in E_{\{2,2\}}} m_{22}(PD_1) x^2 y^2 \\ &\quad + \sum_{uv \in E_{\{2,3\}}} m_{23}(PD_1) x^2 y^3 \\ &= |E_{\{1,2\}}| xy^2 + |E_{\{1,3\}}| x^1 y^3 \\ &\quad + |E_{\{2,2\}}| x^2 y^2 + |E_{\{2,3\}}| x^2 y^3 \\ &= 3 \times 2^n xy^2 + (6 \times 2^n - 3) xy^3 \\ &\quad + (18 \times 2^n - 9) x^2 y^2 \\ &\quad + (21 \times 2^n - 12) x^2 y^3 \\ &= 3 \times 2^n xy^2 + 3(2^{n+1} - 1) xy^3 \\ &\quad + 9(2^{n+1} - 1) x^2 y^2 \\ &\quad + 3(7 \times 2^n - 4) x^2 y^3. \end{aligned} \quad (16)$$

□

Theorem 3. For the PAMAM dendrimers PD_1 , we have

1. $M_1(G) = 105 \times 2^{n+1} - 108.$
2. $M_2(G) = 111 \times 2^{n+1} - 117.$
3. ${}^m M_2(G) = 2^{n+1} + 19 \times 2^{n-1} - 21/4.$
4. $R_\alpha(G) = 3 \times 2^{n+\alpha} + (3^{\alpha+1} + 2^{2\alpha} \times 9)(2^{n+1} - 1) + 2^\alpha \times 3^{\alpha+1}(7 \cdot 2^n - 4).$
5. $R_\alpha(G) = 3 \times 2^{n-\alpha} + (1/3^{\alpha-1} + 9/2^{2\alpha})(2^{n+1} - 1) + (1/(3^{\alpha-1} \times 2^\alpha))(7 \cdot 2^n - 4).$
6. $SSD(G) = 7 \times 2^{n+3} + 3 \times 2^{n+1} + 47 \times 2^n - 54.$
7. $H(G) = (7/5) \times 2^{n+4} - 54/5$
8. $I(G) = (497/20) \times 2^{n+1} - 513/20.$
9. $A(G) = 3 \times 2^{n+3} + 21 \times 2^{n+2} + 369 \times 2^{n-2} - 753/8.$

Proof. Let PD_1 denote PAMAM dendrimers with trifunctional core unit generated by dendrimer generations G_n with n growth stages. Let

$$\begin{aligned} M(G; x, y) &= f(x, y) \\ &= 3 \times 2^n xy^2 + 3(2^{n+1} - 1) xy^3 \\ &\quad + 9(2^{n+1} - 1) x^2 y^2 \\ &\quad + 3(7 \times 2^n - 4) x^2 y^3. \end{aligned} \quad (17)$$

Then

$$\begin{aligned} D_x f(x, y) &= 3 \times 2^n x y^2 + 3(2^{n+1} - 1) x y^3 \\ &\quad + 18(2^{n+1} - 1) x^2 y^2 + 6(7 \times 2^n - 4) x^2 y^3. \end{aligned}$$

$$\begin{aligned} D_y f(x, y) &= 3 \times 2^{n+1} x y^2 + 9(2^{n+1} - 1) x y^3 \\ &\quad + 18(2^{n+1} - 1) x^2 y^2 + 9(7 \times 2^n - 4) x^2 y^3, \end{aligned}$$

$$\begin{aligned} D_y D_x f(x, y) &= 3 \times 2^{n+1} x y^2 + 9(2^{n+1} - 1) x y^3 \\ &\quad + 36(2^{n+1} - 1) x^2 y^2 + 18(7 \times 2^n - 4) x^2 y^3, \end{aligned}$$

$$\begin{aligned} S_x S_y(f(x, y)) &= 3 \times 2^{n-1} x y^2 + (2^{n+1} - 1) x y^3 \\ &\quad + \frac{9}{4}(2^{n+1} - 1) x^2 y^2 + \frac{1}{2}(7 \times 2^n - 4) x^2 y^3, \end{aligned}$$

$$\begin{aligned} D_x^\alpha D_y^\alpha(f(x, y)) &= 3 \times 2^{n+\alpha} x y^2 + 3^{\alpha+1}(2^{n+1} - 1) x y^3 + 2^{2\alpha} \\ &\quad \times 9(2^{n+1} - 1) x^2 y^2 + 2^\alpha \times 3^\alpha (7 \times 2^n - 4) x^2 y^3, \end{aligned}$$

$$\begin{aligned} S_x^\alpha S_y^\alpha(f(x, y)) &= 3 \times 2^{n-\alpha} x y^2 + \frac{1}{3^{\alpha-1}}(2^{n+1} - 1) x y^3 \\ &\quad + \frac{9}{2^{2\alpha}}(2^{n+1} - 1) x^2 y^2 \\ &\quad + \frac{1}{3^{\alpha-1} \times 2^\alpha} (7 \times 2^n - 4) x^2 y^3, \end{aligned}$$

$$\begin{aligned} S_y D_x(f(x, y)) &= 3 \times 2^{n-1} x y^2 + (2^{n+1} - 1) x y^3 \\ &\quad + 9(2^{n+1} - 1) x^2 y^2 + 2(7 \times 2^n - 4) x^2 y^3, \end{aligned}$$

$$\begin{aligned} S_x D_y(f(x, y)) &= 3 \times 2^{n-1} x y^2 + 9(2^{n+1} - 1) x y^3 \\ &\quad + 9(2^{n+1} - 1) x^2 y^2 + \frac{9}{2}(7 \times 2^n - 4) x^2 y^3, \end{aligned}$$

$$\begin{aligned} S_x J f(x, y) &= 2^n x^3 + 3(2^{n+1} - 1) x^4 + \frac{3}{5}(7 \times 2^n - 4) x^5, \end{aligned}$$

$$\begin{aligned} S_x J D_x D_y f(x, y) &= 2^{n+1} x^3 + \frac{45}{4}(2^{n+1} - 1) x^4 + \frac{18}{5}(7 \times 2^n - 4) x^5, \end{aligned}$$

$$\begin{aligned} S_x^3 Q_{-2} J D_x^3 D_y^3 f(x, y) &= 3 \times 2^{n+3} x + \frac{369}{8}(2^{n+1} - 1) x^2 \\ &\quad + 12(7 \times 2^n - 4) x^3. \end{aligned} \tag{18}$$

□

(1) First Zagreb Index

$$\begin{aligned} M_1(G) &= (D_x + D_y) f(x, y) \Big|_{x=y=1} \\ &= 105 \times 2^{n+1} - 108. \end{aligned} \tag{19}$$

(2) Second Zagreb Index

$$M_2(G) = D_y D_x(f(x, y)) \Big|_{x=y=1} = 111 \times 2^{n+1} - 117. \tag{20}$$

(3) Modified Second Zagreb Index

$$\begin{aligned} {}^m M_2(G) &= S_x S_y(f(x, y)) \Big|_{x=y=1} \\ &= 2^{n+1} + 19 \times 2^{n-1} - \frac{21}{4}. \end{aligned} \tag{21}$$

(4) Generalized Randić Index

$$\begin{aligned} R_\alpha(G) &= D_x^\alpha D_y^\alpha(f(x, y)) \Big|_{x=y=1} \\ &= 3 \times 2^{n+\alpha} + (3^{\alpha+1} + 2^{2\alpha} \times 9)(2^{n+1} - 1) + 2^\alpha \\ &\quad \times 3^{\alpha+1} (7 \cdot 2^n - 4). \end{aligned} \tag{22}$$

(5) Inverse Randić Index

$$\begin{aligned} RR_\alpha(G) &= S_x^\alpha S_y^\alpha(f(x, y)) \Big|_{x=y=1} \\ &= 3 \times 2^{n-\alpha} + \left(\frac{1}{3^{\alpha-1}} + \frac{9}{2^{2\alpha}} \right) (2^{n+1} - 1) \\ &\quad + \frac{1}{3^{\alpha-1} \times 2^\alpha} (7 \cdot 2^n - 4). \end{aligned} \tag{23}$$

(6) Symmetric Division Index

$$\begin{aligned}
 SSD(G) &= (S_y D_x + S_x D_y)(f(x, y))\Big|_{x=y=1} \\
 &= 7 \times 2^{n+3} + 3 \times 2^{n+1} + 47 \times 2^n - 54.
 \end{aligned}
 \tag{24}$$

(7) Harmonic Index

$$\begin{aligned}
 H(G) &= 2S_x J(f(x, y))\Big|_{x=1} = H(G) \\
 &= \frac{7}{5} \times 2^{n+4} - \frac{54}{5}.
 \end{aligned}
 \tag{25}$$

(8) Inverse Sum Index

$$I(G) = S_x J D_x D_y (f(x, y))_{x=1} = \frac{497}{20} \times 2^{n+1} - \frac{513}{20}. \tag{26}$$

(9) Augmented Zagreb Index

Theorem 4. For the PAMAM dendrimers PD_2 , we have

$$\begin{aligned}
 M(PD_2, x, y) &= 2^{n+2}xy^2 + 4(2^{n+1} - 1)xy^3 \\
 &\quad + (24 \cdot 2^n - 11)x^2y^2 \\
 &\quad + 14(2^{n+1} - 1)x^2y^3.
 \end{aligned}
 \tag{27}$$

Proof. Let PD_2 be the PAMAM dendrimers with different core generated by dendrimer generators G_n with n growth stages. Then the edge set of PD_2 has following four partitions:

$$\begin{aligned}
 E_{\{1,2\}} &= \{e = uv \in E(PD_2) \mid d_u = 1, d_v = 2\}, \\
 E_{\{1,3\}} &= \{e = uv \in E(PD_2) \mid d_u = 1, d_v = 3\}, \\
 E_{\{2,2\}} &= \{e = uv \in E(PD_2) \mid d_u = 2, d_v = 2\}, \\
 E_{\{2,3\}} &= \{e = uv \in E(PD_2) \mid d_u = 2, d_v = 3\}.
 \end{aligned}
 \tag{28}$$

Now

$$\begin{aligned}
 |E_{\{1,2\}}| &= 4 \cdot 2^n, \\
 |E_{\{1,3\}}| &= 8 \cdot 2^n - 4, \\
 |E_{\{2,2\}}| &= 24 \cdot 2^n - 11,
 \end{aligned}
 \tag{29}$$

and

$$\begin{aligned}
 |E_{\{2,3\}}| &= 28 \cdot 2^n - 14. \\
 M(PD_2; x, y) &= \sum_{i \leq j} m_{ij}(PD_2) x^i y^j \\
 &= \sum_{1 \leq 2} m_{12}(PD_2) xy^2 \\
 &\quad + \sum_{1 \leq 3} m_{13}(PD_2) x^1 y^3 \\
 &\quad + \sum_{2 \leq 2} m_{22}(PD_2) x^2 y^2 \\
 &\quad + \sum_{2 \leq 3} m_{23}(PD_2) x^2 y^3 \\
 &= \sum_{uv \in E_{\{1,2\}}} m_{12}(PD_2) xy^2 \\
 &\quad + \sum_{uv \in E_{\{1,3\}}} m_{13}(PD_2) xy^3 \\
 &\quad + \sum_{uv \in E_{\{2,2\}}} m_{22}(PD_2) x^2 y^2 \\
 &\quad + \sum_{uv \in E_{\{2,3\}}} m_{23}(PD_2) x^2 y^3 \\
 &= |E_{\{1,2\}}| xy^2 + |E_{\{1,3\}}| x^1 y^3 \\
 &\quad + |E_{\{2,2\}}| x^2 y^2 + |E_{\{2,3\}}| x^2 y^3 \\
 &= 4 \cdot 2^n xy^2 + (8 \cdot 2^n - 4) xy^3 \\
 &\quad + (24 \cdot 2^n - 11) x^2 y^2 \\
 &\quad + (28 \cdot 2^n - 14) x^2 y^3 \\
 &= 2^{n+2} xy^2 + 4(2^{n+1} - 1) xy^3 \\
 &\quad + (24 \cdot 2^n - 11) x^2 y^2 \\
 &\quad + 14(2^{n+1} - 1) x^2 y^3.
 \end{aligned}
 \tag{30}$$

□

Theorem 5. For the PAMAM dendrimers PD_2 , we have

1. $M_1(G) = 5(7 \times 2^{n+3} - 26)$.
2. $M_2(G) = 37 \times 2^{n+3} - 140$.
3. ${}^m M_2(G) = (23/3) \times 2^{n+1} - 77/12$.
4. $R_\alpha(G) = 2^{n+\alpha+2} + (4 \times 3^\alpha + 2^{\alpha+1} \times 3^\alpha \times 7)(2^{n+1} - 1) + 2^{2\alpha}(24 \cdot 2^n - 11)$.
5. $R_\alpha(G) = 2^n + (4/3^\alpha + 14/6^\alpha)(2^{n+1} - 1) + (1/2^{2\alpha})(24 \cdot 2^n - 11)$.
6. $SSD(G) = (109/3) \times 2^{n+2} - 197/3$.
7. $H(G) = (7/5) \times 2^{n+6} - 131/10$.

$$8. I(G) = (497/15) \times 2^{n+1} - 154/5.$$

$$9. A(G) = 475 \times 2^n - 427/2.$$

Theorem 6. For the PAMAM dendrimers DS_1 , we have

$$M(DS_1; x, y) = 4 \cdot 3^n xy^4 + 10(3^n - 1)x^2 y^2 + 4(3^n - 1)x^2 y^4. \quad (31)$$

Proof. Let DS_1 be kinds of PAMAM dendrimers with n growth stages.

The edge set of DS_1 has the following three partitions:

$$\begin{aligned} E_{\{1,4\}} &= \{e = uv \in E(DS_1) \mid d_u = 1, d_v = 4\}, \\ E_{\{2,2\}} &= \{e = uv \in E(DS_1) \mid d_u = 2, d_v = 2\}, \\ E_{\{2,4\}} &= \{e = uv \in E(DS_1) \mid d_u = 2, d_v = 4\}. \end{aligned} \quad (32)$$

Now

$$\begin{aligned} |E_{\{1,4\}}| &= 4 \cdot 3^n, \\ |E_{\{2,2\}}| &= 10 \cdot 3^n - 10, \end{aligned} \quad (33)$$

and

$$\begin{aligned} |E_{\{2,4\}}| &= 4 \cdot 3^n - 4. \\ M(DS_1; x, y) &= \sum_{i \leq j} m_{ij}(DS_1) x^i y^j \\ &= \sum_{1 \leq 4} m_{14}(DS_1) xy^4 \\ &\quad + \sum_{2 \leq 2} m_{22}(DS_1) x^2 y^2 \\ &\quad + \sum_{2 \leq 4} m_{24}(DS_1) x^2 y^4 \\ &= \sum_{uv \in E_{\{1,4\}}} m_{14}(DS_1) xy^4 \\ &\quad + \sum_{uv \in E_{\{2,2\}}} m_{22}(DS_1) x^2 y^2 \\ &\quad + \sum_{uv \in E_{\{2,4\}}} m_{24}(DS_1) x^2 y^4 \\ &= |E_{\{1,4\}}| xy^4 + |E_{\{2,2\}}| x^2 y^2 \\ &\quad + |E_{\{2,4\}}| x^2 y^4 \\ &= 4 \cdot 3^n xy^4 + (10 \cdot 3^n - 10) x^2 y^2 \\ &\quad + (4 \cdot 3^n - 4) x^2 y^4 \\ &= 4 \cdot 3^n xy^4 + 10(3^n - 1) x^2 y^2 \\ &\quad + 4(3^n - 1) x^2 y^4. \end{aligned} \quad (34)$$

Theorem 7. For the PAMAM dendrimers DS_1 , we have

1. $M_1(G) = 4(7 \times 3^{n+1} - 16).$
2. $M_2(G) = 8(11 \times 3^n - 9).$
3. ${}^m M_2(G) = 4 \times 3^n - 3.$
4. $R_\alpha(G) = 3^n \times 4^{\alpha+1} + (2^{\alpha+1} \times 5 + 2^{3\alpha+2})(3^n - 1).$
5. $R_\alpha(G) = 3^n/4^{\alpha-1} + (5/2^{2\alpha-1} + 1/2^{3\alpha-2})(3^n - 1).$
6. $SSD(G) = 47 \times 3^n - 30$
7. $H(G) = (119/15) \times 3^n - 19/5$
8. $I(G) = (278/15) \times 3^n - 46/3.$
9. $A(G) = (3280/27) \times 3^n - 112.$

3.2. M-Polynomials and Degree-Based Indices for Polyomino Chains. From the geometric point of view, a polyomino system is a finite 2-connected plane graph in which each interior cell is encircled by a regular square. In other words, it is an edge-connected union of cells in the planar square lattice. Polyomino chain is a particular polyomino system such that the joining of the centers (set c_i as the center of the i th square) of its adjacent regular composes a path $c_1, c_2, c_3, \dots, c_n$.

Let B_n be the set of polyomino chains with n squares. There are $2n+1$ edges in every $B_n \in B_n$, where B_n is named as a linear chain and denoted by L_n if the subgraph of B_n induced by the vertices with $d(v)=3$ is a molecular graph with exactly $n-2$ squares. Also, B_n can be called a zigzag chain and labelled as Z_n if the subgraph of B_n is induced by the vertices with $d(v)>2$ is P_n .

The angularly connected, or branched, squares constitute a link of a polyomino chain. A maximal linear chain (containing the terminal squares and kinks at its end) in the polyomino chains is called a segment of polyomino chain. Let $l(S)$ be the length of S which is calculated by the number of squares in S . For any segment S of a polyomino chain, we get $l(S) \in \{2, 3, 4, \dots, n\}$. Furthermore, we deduce $l_1 = n$ and $m=1$ for a linear chain L_n with n squares and $l_i = 2$ and $m=n-1$ for a zigzag chain Z_n with n squares.

In what follows, we always assume that a polyomino chain consists of a sequence of segments $S_1, S_2, S_3, \dots, S_m$ and $L(S_i) = l_i$, where $m \geq 1$ and $i \in \{2, 3, 4, \dots, m\}$. We derive that $\sum_{i=1}^m l_i = n + m - 1$.

Theorem 8. For a linear polyomino chain L_n , we have $M(L_n; x, y) = 2x^2 y^2 + 4x^2 y^3 + (3n - 5)x^3 y^3$.

Proof. Let L_n be the polyomino chain with n squares where $l_1 = n$ and $m=1$. L_n is called the linear chain.

The edge set of L_n has the following three partitions:

$$\begin{aligned} E_{\{2,2\}} &= \{e = uv \in E(L_n) \mid d_u = 2, d_v = 2\}, \\ E_{\{2,3\}} &= \{e = uv \in E(L_n) \mid d_u = 2, d_v = 3\}, \\ E_{\{3,3\}} &= \{e = uv \in E(L_n) \mid d_u = 3, d_v = 3\}. \end{aligned} \quad (35)$$

Now

$$|E_{\{2,2\}}| = 2, \quad (36)$$

□

and

$$|E_{\{2,3\}}| = 4, \tag{37}$$

$$|E_{\{3,3\}}| = 3n - 5.$$

$$\begin{aligned} M(L_n; x, y) &= \sum_{i \leq j} m_{ij}(L_n) x^i y^j \\ &= \sum_{2 \leq 2} m_{22}(L_n) x^2 y^2 + \sum_{2 \leq 3} m_{23}(L_n) x^2 y^3 \\ &\quad + \sum_{3 \leq 3} m_{33}(L_n) x^3 y^3 \\ &= \sum_{uv \in E_{\{2,2\}}} m_{22}(L_n) x^2 y^2 \\ &\quad + \sum_{uv \in E_{\{2,3\}}} m_{23}(L_n) x^2 y^3 \\ &\quad + \sum_{uv \in E_{\{3,3\}}} m_{33}(L_n) x^3 y^3 \\ &= |E_{\{2,2\}}| x^2 y^2 + |E_{\{2,3\}}| x^2 y^3 \\ &\quad + |E_{\{3,3\}}| x^3 y^3 \\ &= 2x^2 y^2 + 4x^2 y^3 + (3n - 5) x^3 y^3. \end{aligned} \tag{38}$$

□

Theorem 9. For a linear polyomino chain L_n , we have the following:

1. $M_1(G) = 18n - 2.$
2. $M_2(G) = 27n - 13.$
3. ${}^m M_2(G) = (1/3)n - 11/18.$
4. $R_\alpha(G) = 2^{2\alpha+1} + 2^{\alpha+2} \cdot 3^\alpha + 3^{2\alpha}(3n - 5).$
5. $R_\alpha(G) = 1/2^{\alpha-1} + 2^{2-\alpha}/3^\alpha + (1/3^{2\alpha})(3n - 5).$
6. $SSD(G) = 6n + 8/3.$
7. $H(G) = n + 14/15.$
8. $I(G) = (9/2)n - 7/10.$
9. $A(G) = (2187/64)n - 573/64.$

Theorem 10. Let Z_n be zigzag polyomino chain with n squares such that $l_i = 2$ and $m = n - 1$. Then

$$\begin{aligned} M(Z_n; x, y) &= 2x^2 y^2 + 4x^2 y^3 + 2(m - 1) x^2 y^4 \\ &\quad + 2x^3 y^4 + (3n - 2m - 5) x^4 y^4. \end{aligned} \tag{39}$$

Proof. Let Z_n be zigzag polyomino chain with n squares such that $l_i = 2$ and $m = n - 1$. Polyomino chain consists of a sequence of segments S_1, S_2, \dots, S_m and $l(S_i) = l_i$ where $m \geq 1$ and $i \in \{1, 2, \dots, m\}$.

The edge set of Z_n has the following five partitions:

$$\begin{aligned} E_{\{2,2\}} &= \{e = uv \in E(Z_n) \mid d_u = 2, d_v = 2\}, \\ E_{\{2,3\}} &= \{e = uv \in E(Z_n) \mid d_u = 2, d_v = 3\}, \\ E_{\{2,4\}} &= \{e = uv \in E(Z_n) \mid d_u = 2, d_v = 4\}, \\ E_{\{3,4\}} &= \{e = uv \in E(Z_n) \mid d_u = 3, d_v = 4\}, \\ E_{\{4,4\}} &= \{e = uv \in E(Z_n) \mid d_u = 4, d_v = 4\}. \end{aligned} \tag{40}$$

Now

$$\begin{aligned} |E_{\{2,2\}}| &= 2, \\ |E_{\{2,3\}}| &= 4, \\ |E_{\{2,4\}}| &= 2(m - 1), \\ |E_{\{3,4\}}| &= 2, \end{aligned} \tag{41}$$

and

$$\begin{aligned} |E_{\{4,4\}}| &= 3n - 2m - 5. \\ M(Z_n; x, y) &= \sum_{i \leq j} m_{ij}(Z_n) x^i y^j \\ &= \sum_{2 \leq 2} m_{22}(Z_n) x^2 y^2 + \sum_{2 \leq 3} m_{23}(Z_n) x^2 y^3 \\ &\quad + \sum_{2 \leq 4} m_{24}(Z_n) x^2 y^4 \\ &\quad + \sum_{3 \leq 4} m_{34}(Z_n) x^3 y^4 \\ &\quad + \sum_{4 \leq 4} m_{44}(Z_n) x^4 y^4 \\ &= \sum_{uv \in E_{\{2,2\}}} m_{22}(Z_n) x^2 y^2 \\ &\quad + \sum_{uv \in E_{\{2,3\}}} m_{23}(Z_n) x^2 y^3 \\ &\quad + \sum_{uv \in E_{\{2,4\}}} m_{24}(Z_n) x^2 y^4 \\ &\quad + \sum_{uv \in E_{\{3,4\}}} m_{34}(Z_n) x^3 y^4 \\ &\quad + \sum_{uv \in E_{\{4,4\}}} m_{44}(Z_n) x^4 y^4 \\ &= |E_{\{2,2\}}| x^2 y^2 + |E_{\{2,3\}}| x^2 y^3 \\ &\quad + |E_{\{2,4\}}| x^2 y^4 + |E_{\{3,4\}}| x^3 y^4 \\ &\quad + |E_{\{4,4\}}| x^4 y^4 \\ &= 2x^2 y^2 + 4x^2 y^3 + 2(m - 1) x^2 y^4 \\ &\quad + 2x^3 y^4 + (3n - 2m - 5) x^4 y^4. \end{aligned} \tag{42}$$

□

Theorem 11. For the Zigzag polyomino chain Z_n for $n \geq 2$, we have the following:

1. $M_1(G) = 24n - 4m - 10$.
2. $M_2(G) = 48n - 16m - 40$.
3. ${}^m M_2(G) = (1/2)n + (1/3)m + 37/18$
4. $R_\alpha(G) = 2^{2\alpha+1} + 2^{\alpha+2} \times 3^\alpha + 2^{3\alpha+1}(m-1) + 3^\alpha \times 2^{2\alpha+1} + 2^{4\alpha}(3n - 2m - 5)$.
5. $R_\alpha(G) = 1/2^{2\alpha-1} + 1/(3^\alpha \cdot 2^{\alpha-2}) + (1/2^{3\alpha-1})(m-1) + 1/(2^{2\alpha-1} \times 3^\alpha) + (1/2^{4\alpha})(3n - 2m - 5)$.
6. $SSD(G) = 6n + m + 11/6$
7. $H(G) = (3/4)n + (1/6)m + 61/60$.
8. $I(G) = (3/5)n - (4/3)m - 92/35$.
9. $A(G) = (512/9)n - (592/27)m - 118688/3375$.

Theorem 12. For the polyomino chain with n squares and of m segments S_1 and S_2 satisfying $l_1 = 2$ and $l_2 = n - 1$, B_n^1 ($n \geq 3$), we have

$$M(B_n^1, x, y) = 2x^2y^2 + 5x^2y^3 + x^2y^4 + (3n - 10)x^3y^3 + 3x^3y^4. \quad (43)$$

Proof. Let B_n^1 ($n \geq 3$) be the polyomino chain with n squares and of m segments S_1 and S_2 satisfying $l_1 = 2$ and $l_2 = n - 1$. The edge set of B_n^1 ($n \geq 3$) has the following five partitions:

$$\begin{aligned} E_{\{2,2\}} &= \{e = uv \in E(B_n^1) \mid d_u = 2, d_v = 2\}, \\ E_{\{2,3\}} &= \{e = uv \in E(B_n^1) \mid d_u = 2, d_v = 3\}, \\ E_{\{2,4\}} &= \{e = uv \in E(B_n^1) \mid d_u = 2, d_v = 4\}, \\ E_{\{3,3\}} &= \{e = uv \in E(B_n^1) \mid d_u = 3, d_v = 3\}, \\ E_{\{3,4\}} &= \{e = uv \in E(B_n^1) \mid d_u = 3, d_v = 4\}. \end{aligned} \quad (44)$$

Now

$$\begin{aligned} |E_{\{2,2\}}| &= 2, \\ |E_{\{2,3\}}| &= 5, \\ |E_{\{2,4\}}| &= 1, \\ |E_{\{3,3\}}| &= 3n - 10, \end{aligned} \quad (45)$$

and

$$|E_{\{3,4\}}| = 3. \quad (46)$$

$$\begin{aligned} M(B_n^1; x, y) &= \sum_{i \leq j} m_{ij}(B_n^1) x^i y^j \\ &= \sum_{2 \leq 2} m_{22}(B_n^1) x^2 y^2 + \sum_{2 \leq 3} m_{23}(B_n^1) x^2 y^3 \\ &\quad + \sum_{2 \leq 4} m_{24}(B_n^1) x^2 y^4 \\ &\quad + \sum_{3 \leq 3} m_{33}(B_n^1) x^3 y^3 \\ &\quad + \sum_{3 \leq 4} m_{34}(B_n^1) x^3 y^4 \\ &= \sum_{uv \in E_{\{2,2\}}} m_{22}(B_n^1) x^2 y^2 \\ &\quad + \sum_{uv \in E_{\{2,3\}}} m_{23}(B_n^1) x^2 y^3 \\ &\quad + \sum_{uv \in E_{\{2,4\}}} m_{24}(B_n^1) x^2 y^4 \\ &\quad + \sum_{uv \in E_{\{3,3\}}} m_{33}(B_n^1) x^3 y^3 \\ &\quad + \sum_{uv \in E_{\{3,4\}}} m_{34}(B_n^1) x^3 y^4 \\ &= |E_{\{2,2\}}| x^2 y^2 + |E_{\{2,3\}}| x^2 y^3 \\ &\quad + |E_{\{2,4\}}| x^2 y^4 + |E_{\{3,3\}}| x^3 y^3 \\ &\quad + |E_{\{3,4\}}| x^3 y^4 \\ &= 2x^2 y^2 + 5x^2 y^3 + x^2 y^4 \\ &\quad + (3n - 10)x^3 y^3 + 3x^3 y^4. \end{aligned} \quad (47)$$

□

Theorem 13. For the polyomino chain with n squares and of m segments S_1 and S_2 satisfying $l_1 = 2$ and $l_2 = n - 1$, B_n^1 ($n \geq 3$), we have the following:

1. $M_1(G) = 18n$.
2. $M_2(G) = 27n - 8$.
3. ${}^m M_2(G) = n/3 + 43/72$.
4. $R_\alpha(G) = 2^{2\alpha+1} + 6^\alpha \times 5 + 2^{3\alpha} + 3^{2\alpha}(3n - 10) + 3^{\alpha+1} \times 4^\alpha$.
5. $R_\alpha(G) = 1/2^{2\alpha-1} + 5/6^\alpha + 1/2^{3\alpha} + (1/3^{2\alpha})(3n - 10) + 1/(2^{2\alpha} \times 3^{\alpha-1})$.
6. $SSD(G) = 6n + 43/12$.
7. $H(G) = n + 6/7$.
8. $I(G) = 27n - 709/10$.
9. $A(G) = (2187/64)n - 33737/4000$.

Theorem 14. For polyomino chain with n squares and m segments S_1, S_2, \dots, S_m ($m \geq 3$) satisfying $l_1 = l_m = 2$ and $l_2, \dots, l_{m-1} \geq 3$, B_n^2 ($n \geq 4$), we have

$$\begin{aligned} M(B_n^2, x, y) &= 2x^2y^2 + 2mx^2y^3 + 2x^2y^4 \\ &+ 3(n-2m+1)x^3y^3 \\ &+ 2(2m-3)x^3y^4. \end{aligned} \quad (48)$$

Proof. Let B_n^2 ($n \geq 4$) be a polyomino chain with n squares and m segments S_1, S_2, \dots, S_m ($m \geq 3$) satisfying $l_1 = l_m = 2$ and $l_2, \dots, l_{m-1} \geq 3$. Then the edge set of B_n^2 ($n \geq 4$) has the following five partitions:

$$\begin{aligned} E_{\{2,2\}} &= \{e = uv \in E(B_n^2) \mid d_u = 2, d_v = 2\}, \\ E_{\{2,3\}} &= n \{e = uv \in E(B_n^2) \mid d_u = 2, d_v = 3\}, \\ E_{\{2,4\}} &= \{e = uv \in E(B_n^2) \mid d_u = 2, d_v = 4\}, \\ E_{\{3,3\}} &= \{e = uv \in E(B_n^2) \mid d_u = 3, d_v = 3\}, \\ E_{\{3,4\}} &= \{e = uv \in E(B_n^2) \mid d_u = 3, d_v = 4\}. \end{aligned} \quad (49)$$

Now

$$\begin{aligned} |E_{\{2,2\}}| &= 2, \\ |E_{\{2,3\}}| &= 2m, \\ |E_{\{2,4\}}| &= 2, \\ |E_{\{3,3\}}| &= 3n - 6m + 3, \end{aligned} \quad (50)$$

and

$$\begin{aligned} |E_{\{4,4\}}| &= 4m - 6. \\ M(B_n^2; x, y) &= \sum_{i \leq j} m_{ij}(B_n^2) x^i y^j \\ &= \sum_{2 \leq 2} m_{22}(B_n^2) x^2 y^2 + \sum_{2 \leq 3} m_{23}(B_n^2) x^2 y^3 \\ &+ \sum_{2 \leq 4} m_{24}(B_n^2) x^2 y^4 + \sum_{3 \leq 3} m_{33}(B_n^2) x^3 y^3 \\ &+ \sum_{3 \leq 4} m_{34}(B_n^2) x^3 y^4 \\ &= \sum_{uv \in E_{\{2,2\}}} m_{22}(B_n^2) x^2 y^2 + \sum_{uv \in E_{\{2,3\}}} m_{23}(B_n^2) x^2 y^3 \\ &+ \sum_{uv \in E_{\{2,4\}}} m_{24}(B_n^2) x^2 y^4 + \sum_{uv \in E_{\{3,3\}}} m_{33}(B_n^2) x^3 y^3 \end{aligned} \quad (51)$$

$$\begin{aligned} &+ \sum_{uv \in E_{\{3,4\}}} m_{34}(B_n^2) x^3 y^4 \\ &= |E_{\{2,2\}}| x^2 y^2 + |E_{\{2,3\}}| x^2 y^3 + |E_{\{2,4\}}| x^2 y^4 \\ &+ |E_{\{3,3\}}| x^3 y^3 + |E_{\{3,4\}}| x^3 y^4 \\ &= 2x^2 y^2 + 2mx^2 y^3 + 2x^2 y^4 + (3n - 6m + 3) x^3 y^3 \\ &+ (4m - 6) x^3 y^4 \\ &= 2x^2 y^2 + 2mx^2 y^3 + 2x^2 y^4 + 3(n - 2m + 1) x^3 y^3 \\ &+ 2(2m - 3) x^3 y^4 \end{aligned} \quad (52)$$

□

Theorem 15. For polyomino chain with n squares and m segments S_1, S_2, \dots, S_m ($m \geq 3$) satisfying $l_1 = l_m = 2$ and $l_2, \dots, l_{m-1} \geq 3$, B_n^2 ($n \geq 4$), we have the following:

1. $M_1(G) = 2(9n + m - 2)$.
2. $M_2(G) = 3(9n + 2m - 7)$.
3. ${}^m M_2(G) = (1/3)n + 7/12$
4. $R_\alpha(G) = 3^{2\alpha+1}n + (3^\alpha \times 2^{\alpha+1} - 2 \times 3^{2\alpha+1} + 3^\alpha \times 2^{2\alpha+2})m + (1 + 3^{\alpha+1})2^{2\alpha+1} + 2^{3\alpha+1} + 3^{2\alpha+1}$.
5. $R_\alpha(G) = (1/3^{2\alpha-1})n + (1/(3^\alpha \cdot 2^{\alpha-1}) + 2/3^{2\alpha-1} + 1/(2^{2\alpha-2} \times 3^\alpha))m + (1 - 1/3^{\alpha-1})(1/2^{2\alpha-1}) + 1/2^{3\alpha-1} + 1/3^{2\alpha-1}$.
6. $SSD(G) = 6n + (2/3)m + 5/2$
7. $H(G) = n - (2/35)m + 20/21$.
8. $I(G) = (9/2)n + (9/35)m - 47/42$.
9. $A(G) = (2187/64)n + (11809/4000)m - 134177/8000$.

4. Conclusions

Topological indices calculated in this paper help us to guess biological activities, chemical reactivity, and physical features of under-study dendrimers and polyomino chains. For example, Randić index is useful for determining physicochemical properties of alkanes as noticed by chemist Milan Randić in 1975. He noticed the correlation between the Randić index and several physicochemical properties of alkanes like boiling point, vapor pressure, enthalpies of formation, surface area, and chromatographic retention times. Hence our results are helpful in determination of the significance of PAMAM dendrimers and polyomino chains in pharmacy and industry.

Data Availability

All data required for this research is included within this paper.

Conflicts of Interest

The authors do not have any conflicts of interest.

Authors' Contributions

All authors contributed equally to this paper.

Acknowledgments

The first author is funded by Dong-A University, Korea.

References

- [1] T. Go, *Handbook of discrete and computational geometry*, J. E. Goodman and J. O'Rourke, Eds., CRC Press Series on Discrete Mathematics and its Applications, CRC Press, Boca Raton, FL, 1997.
- [2] S. W. Golomb, *Polyominoes, Puzzles, Patterns, Problems, And Packings*, Princeton University Press, 1996.
- [3] L. Alonso and R. Cerf, "The three dimensional polyominoes of minimal area," *Electronic Journal of Combinatorics*, vol. 3, no. 1 R, pp. 1-39, 1996.
- [4] D. Astruc, E. Boisselier, and C. Ornelas, "Dendrimers designed for functions: from physical, photophysical, and supramolecular properties to applications in sensing, catalysis, molecular electronics, photonics, and nanomedicine," *Chemical Reviews*, vol. 110, no. 4, pp. 1857-1959, 2010.
- [5] F. Vögtle, G. Richardt, and N. Werner, *Dendrimer Chemistry Concepts, Syntheses, Properties, Applications*, 2009.
- [6] M. R. Farahani, W. Gao, and M. R. Kanna, "The connective eccentric index for an infinite family of dendrimers," *Journal of Fundamental and Applied Life Sciences*, vol. 5, no. S4, pp. 766-771, 2015.
- [7] W. Gao, L. Shi, and M. R. Farahani, "Distance-based indices for some families of dendrimer nanostars," *IAENG International Journal of Applied Mathematics*, vol. 46, no. 2, pp. 168-186, 2016.
- [8] W. Gao and W. Wang, "Degree-based indices of polyhex nanotubes and dendrimer nanostar," *Journal of Computational and Theoretical Nanoscience*, vol. 13, no. 3, pp. 1577-1583, 2016.
- [9] W. Gao and M. R. Farahani, "The hyper-zagreb index for an infinite family of nanostar dendrimer," *Journal of Discrete Mathematical Sciences & Cryptography*, vol. 20, no. 2, pp. 515-523, 2017.
- [10] W. Gao, M. R. Farahani, and L. Shi, "The forgotten topological index of some drug structures," *Acta Medica Mediterranea*, vol. 32, no. 1, pp. 579-585, 2016.
- [11] W. Gao, Y. Wang, W. Wang, and L. Shi, "The first multiplication atom-bond connectivity index of molecular structures in drugs," *Saudi Pharmaceutical Journal*, vol. 25, no. 4, pp. 548-555, 2017.
- [12] W. Gao, M. Younas, A. Farooq et al., "M-Polynomials and Degree-Based Topological Indices of the Crystallographic Structure of Molecules," *Biomolecules*, vol. 8, no. 4, p. 107, 2018.
- [13] H. Zhang and F. Zhang, "The Clar covering polynomial of hexagonal systems," *Discrete Applied Mathematics*, vol. 69, no. 1-2, pp. 147-167, 1996.
- [14] H. Wiener, "Structural determination of paraffin boiling points," *Journal of the American Chemical Society*, vol. 69, no. 1, pp. 17-20, 1947.
- [15] E. Deutsch and S. Klavžar, "M-Polynomial and degree-based topological indices," *Iranian Journal of Mathematical Chemistry*, vol. 6, pp. 93-102, 2015.
- [16] M. Munir, W. Nazeer, S. Rafique, and S. M. Kang, "M-polynomial and related topological indices of nanostar dendrimers," *Symmetry*, vol. 8, no. 9, p. 97, 2016.
- [17] M. Munir, W. Nazeer, A. R. Nizami, S. Rafique, and S. M. Kang, "M-polynomials and topological indices of titania nanotubes," *Symmetry*, vol. 8, no. 11, p. 117, 2016.
- [18] Y. C. Kwun, M. Munir, W. Nazeer, S. Rafique, and S. Min Kang, "M-Polynomials and topological indices of V-Phenylenic Nanotubes and Nanotori," *Scientific Reports*, vol. 7, no. 1, 2017.
- [19] M. Munir, W. Nazeer, S. Rafique, A. R. Nizami, and S. M. Kang, "Some computational aspects of boron triangular nanotubes," *Symmetry*, vol. 9, no. 6, 2017.
- [20] M. Munir, W. Nazeer, S. Rafique, and S. M. Kang, "M-polynomial and degree-based topological indices of polyhex nanotubes," *Symmetry*, vol. 8, no. 12, p. 149, 2016.
- [21] A. A. Dobrynin, R. Entringer, and I. Gutman, "Wiener index of trees: theory and applications," *Acta Applicandae Mathematicae*, vol. 66, no. 3, pp. 211-249, 2001.
- [22] B. Mohar and T. Pisanski, "How to compute the Wiener index of a graph," *Journal of Mathematical Chemistry*, vol. 2, no. 3, pp. 267-277, 1988.
- [23] B. Zhou and I. Gutman, "Relations between Wiener, hyper-Wiener and Zagreb indices," *Chemical Physics Letters*, vol. 394, no. 1-3, pp. 93-95, 2004.
- [24] M. Randić, "On characterization of molecular branching," *Journal of the American Chemical Society*, vol. 97, no. 23, pp. 6609-6615, 1975.
- [25] S. M. Kang, M. A. Zahid, W. Nazeer, and W. Gao, "Calculating the Degree-based Topological Indices of Dendrimers," *Open Chemistry*, vol. 16, no. 1, pp. 681-688, 2018.
- [26] B. Bollobás and P. Erdős, "Graphs of extremal weights," *Ars Combinatoria*, vol. 50, pp. 225-233, 1998.
- [27] D. Amić, D. Bešlo, B. Lučić, S. Nikolić, and N. Trinajstić, "The Vertex-Connectivity Index Revisited," *Journal of Chemical Information and Computer Sciences*, vol. 38, no. 5, pp. 819-822, 1998.
- [28] Y. Hu, X. Li, Y. Shi, T. Xu, and I. Gutman, "On molecular graphs with smallest and greatest zeroth-order general randić index," *MATCH - Communications in Mathematical and in Computer Chemistry*, vol. 54, no. 2, pp. 425-434, 2005.
- [29] G. Caporossi, I. Gutman, P. Hansen, and L. Pavlović, "Graphs with maximum connectivity index," *Computational Biology and Chemistry*, vol. 27, no. 1, pp. 85-90, 2003.
- [30] I. Gutman and K. C. Das, "The first Zagreb index 30 years after," *MATCH Communications in Mathematical and in Computer Chemistry*, vol. 50, no. 1, pp. 83-92, 2004.
- [31] K. C. Das and I. Gutman, "Some properties of the second Zagreb index," *MATCH Communications in Mathematical and in Computer Chemistry*, vol. 52, no. 1, pp. 1-3, 2004.
- [32] I. Gutman, "An exceptional property of first Zagreb index," *MATCH - Communications in Mathematical and in Computer Chemistry*, vol. 72, no. 3, pp. 733-740, 2014.
- [33] Y. Huang, B. Liu, and L. Gan, "Augmented Zagreb index of connected graphs," *MATCH - Communications in Mathematical and in Computer Chemistry*, vol. 67, no. 2, pp. 483-494, 2012.
- [34] W. Nazeer, A. Farooq, M. Younas, M. Munir, and S. Kang, "On Molecular Descriptors of Carbon Nanocones," *Biomolecules*, vol. 8, no. 3, p. 92, 2018.
- [35] Y. Kwun, A. Virk, W. Nazeer, M. Rehman, and S. Kang, "On the Multiplicative Degree-Based Topological Indices of Silicon-Carbon Si₂C₃-I [p, q] and Si₂C₃-II [p, q]," *Symmetry*, vol. 10, no. 8, p. 320, 2018.

- [36] S. Kang, Z. Iqbal, M. Ishaq, R. Sarfraz, A. Aslam, and W. Nazeer, "On Eccentricity-Based Topological Indices and Polynomials of Phosphorus-Containing Dendrimers," *Symmetry*, vol. 10, no. 7, p. 237, 2018.
- [37] Y. C. Kwun, A. Ali, W. Nazeer, M. Ahmad Chaudhary, and S. M. Kang, "M-Polynomials and Degree-Based Topological Indices of Triangular, Hourglass, and Jagged-Rectangle Benzenoid Systems," *Journal of Chemistry*, vol. 2018, Article ID 8213950, 8 pages, 2018.
- [38] J.-B. Liu, X.-F. Pan, L. Yu, and D. Li, "Complete characterization of bicyclic graphs with minimal Kirchhoff index," *Discrete Applied Mathematics*, vol. 200, pp. 95–107, 2016.
- [39] J.-B. Liu, W.-R. Wang, Y.-M. Zhang, and X.-F. Pan, "On degree resistance distance of cacti," *Discrete Applied Mathematics: The Journal of Combinatorial Algorithms, Informatics and Computational Sciences*, vol. 203, pp. 217–225, 2016.
- [40] J.-B. Liu and X.-F. Pan, "Minimizing Kirchhoff index among graphs with a given vertex bipartiteness," *Applied Mathematics and Computation*, vol. 291, pp. 84–88, 2016.
- [41] W. Gao, M. K. Jamil, A. Javed, M. R. Farahani, S. Wang, and J.-B. Liu, "Sharp Bounds of the Hyper-Zagreb Index on Acyclic, Unicyclic, and Bicyclic Graphs," *Discrete Dynamics in Nature and Society*, vol. 2017, Article ID 6079450, 5 pages, 2017.
- [42] J.-B. Liu, W. Gao, M. K. Siddiqui, and M. R. Farahani, "Computing three topological indices for Titania nanotubes $TiO_2[m,n]$," *AKCE International Journal of Graphs and Combinatorics*, vol. 13, no. 3, pp. 255–260, 2016.
- [43] Q. Liu, J.-B. Liu, and J. Cao, "The Laplacian polynomial and Kirchhoff index of graphs based on R-graphs," *Neurocomputing*, vol. 177, pp. 441–446, 2016.
- [44] J.-B. Liu, S. Wang, C. Wang, and S. Hayat, "Further results on computation of topological indices of certain networks," *IET Control Theory & Applications*, vol. 11, no. 13, pp. 2065–2071, 2017.
- [45] Y. Huo, J.-B. Liu, A. Q. Baig, W. Sajjad, and M. R. Farahani, "Connective eccentric index of NaN_m nanotube," *Journal of Computational and Theoretical Nanoscience*, vol. 14, no. 4, pp. 1832–1836, 2017.

