



Characterization and Management of Post-harvest Fungal Diseases in Banana (*Musa paradisiaca*)

Shital Pawar ^{a*}, Gulab Rathod ^a
and Manjunatha Nanjunadappa ^b

^a Department of Botany, Shrikrishna Mahavidyalaya, Gunjoti, Osmanabad-413606, (M.S.), India.

^b ICAR-National Research Centre on Pomegranate, Solapur 413255, (M.S.), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i81230>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119963>

Original Research Article

Received: 25/05/2024

Accepted: 01/08/2024

Published: 07/08/2024

ABSTRACT

Banana (*Musa* spp.) is a vital crop globally, contributing significantly to food security and income generation, particularly in tropical regions like India. However, post-harvest fungal diseases pose a considerable threat to banana production, affecting both fruit quality and marketability. The current study investigated the isolation, characterization, and management of post-harvest fungal diseases in banana (*Musa paradisiaca*). Infected banana fruits were collected from the Dharashiv fruit market (Maharashtra, India), and pathogenic fungi were isolated and identified. *Fusarium napiforme*, *Talaromyces atrovirens*, *Cladosporium cladosporioides*, *Fusarium* sp., and *Fusarium equiseti* were the primary pathogens identified. DNA extraction and sequencing were employed for accurate identification, and sequences were submitted to GenBank. The antifungal activity of essential oils

*Corresponding author: E-mail: pawarshitu333@gmail.com;

and plant extracts was evaluated using the Poisoned Food Technique. Essential oils from *Syzygium aromaticum*, *Mentha piperita*, and *Punica granatum* showed significant inhibition ($P < 0.05$) of fungal growth, with clove and peppermint oils achieving 100% inhibition at higher concentrations. Plant extracts of *Ocimum sanctum*, *Eucalyptus globulus*, *Mentha piperita*, *Zingiber officinale*, *Curcuma longa*, *Azadirachta indica*, *Piper betel*, and *Cymbopogon citratus* were also tested, revealing notable efficacy, particularly with neem and peppermint extracts. The study that the efficacy of essential oils was more compared to aqueous plant extracts. The results suggested sustainable strategies for managing post-harvest fungal diseases in bananas and explained the importance of conducting field trials to validate laboratory results.

Keywords: Banana; ITS; post-harvest; plant extracts; essential oils.

1. INTRODUCTION

Banana (*Musa* spp.) is a perennial plant renowned for its edible fruits, playing a crucial role in food security and income generation for communities worldwide. Bananas are extensively cultivated in subtropical and tropical regions, with annual production estimated at over 102 million tonnes of fresh fruit worldwide with India alone producing 26.5 million tonnes of banana [1,2]. According to Canton [3], global banana production is expected to rise to 140 million tonnes over the next decade, with India projected to remain the leading producer, reaching an output of 35 million tonnes by 2032. However, according to the OECD/FAO [4], there has been a decline in banana production and export, from 20.5 million tonnes in 2021 to 19.6 million tonnes in 2022. One of the important reasons for the decline might be because of the diseases primarily fungal diseases that not only attack the banana plant during its growth stages but also pose a substantial threat to the fruit during the post-harvest period [5,6,7].

The most severe out of all the fungi is *Fusarium oxysporum* f. sp. cubense (Foc) tropical race 4, that has devastated banana fields globally, causing the well-known Panama disease [8]. In June 2015, *Fusarium oxysporum* f. sp. cubense Tropical Race 4 (Foc TR4) was detected in Bihar, India's largest banana-producing province [9]. Previously, Fusarium wilt caused by Foc race 1 affected Cavendish bananas in Theni, southern India [10]. Beyond the field, fungal infections continue to threaten bananas during post-harvest handling and storage. Common post-harvest fungal pathogens include *Burkholderia*, *Pseudomonas*, *Elaphocordyceps*, *Penicillium*, and *Talaromyces* [11]. These infections can significantly reduce fruit quality, marketability, and shelf life, leading to substantial economic losses. Numerous studies have reported post-

harvest fungal diseases affecting bananas across various regions of India [12-16].

Application of essential oils and plant extracts from the plants is another effective way to control post-harvest diseases. Essential oils and plant extracts are known to be safe and would therefore be acceptable among the human population. The oils are known to be biologically active in their vapour phase. In the vapour phase they might act as fumigants and thereby manage post-harvest pathogens. Since the essential oils are more complex with different metabolites, the chances of pathogen resistance to the oils is low [16]. Studies have already been conducted on the efficacy of essential oils from *Cinnamomum zeylanicum*, *Azadirachta indica*, and *Mentha arvensis* that exhibited 100% efficacy against post-harvest banana pathogens [17]. Other alternatives like cinnamon, thyme, and almond oils have been used to protect bananas [18].

The current study focuses on the isolation, characterization, and management of post-harvest fungal diseases in banana by the use of alternative and natural strategies like plant extracts and essential oils.

2. MATERIALS AND METHODS

2.1 Collection and Isolation of the Pathogenic Fungi from the Fruit Samples

100 Infected banana fruits were collected from three different locations within the Dharashiv fruit market (18°11'12.3"N, 76°02'30.4"E). All the 100 fruit samples were selected, sealed in sterile polyethylene bags, and promptly transported to the laboratory for fungal isolation. To obtain pure cultures, infected patches of the peel off fruit samples were used. The isolation process was conducted immediately to minimize the presence of other saprophytic fungi on the fruit surface.

The infected tissue was excised into small pieces of each that contained actively growing conidia were cut from these fruits using a flame sterilised scalpel. The tissues were placed to sterile potato dextrose agar plates (PDA) and incubated at 28 °C for seven days [19]. Petri dishes were observed daily, and the distinct colonies of fungi were picked. The isolated fungi were purified using a single spore technique [20], and the pure colonies of fungal isolates were maintained on PDA slants.

2.2 Identification of Fungal Isolates

DNA extraction followed the method described by Saitoh, Togashi [21]. The Internal Transcribed Spacer (ITS) region of ribosomal DNA was amplified using ITS4(5'-TCCTCCGCTTATTGATATGC -3') and ITS5(5'-GGAAGTAAAAGTCGTAACAAGG -3') primer pairs. The PCR reaction mixture consisted of 2.5 µl of 10X PCR buffer, 1 µl of 200 mM dNTPs, 0.2 µl of Taq polymerase (1U/µl), 1 µl each of 10 pM/µl ITS5 and ITS4 primers, and 2.5 µl of 10 ng/µl template DNA. Sequencing of the amplified product was performed using an Applied Biosystems Sanger sequencer (ABI 3100 Avant Prism, United States). Consensus sequences were obtained from forward and reverse complementary sequences, and sequence identification was conducted using nBLAST alignment. Phylogenetic analysis for identifying the isolates was performed using MEGA 11 [22]. The identified pathogenic fungal sequences were submitted to GenBank. The phylogenetic analysis of ITS region of the fungal samples are shown from Figs. 1 to 5.

2.3 In-vitro Antifungal Activity Using Poisoned Food Technique

Fresh plant materials including *Ocimum sanctum*, *Eucalyptus globulus*, *Mentha piperita*, *Zingiber officinale*, *Curcuma longa*, *Azadirachta indica*, *Piper betel*, and *Cymbopogon citratus* were collected. The leaves were washed with sterilized distilled water, shade-dried for 5 days, and ground into a fine powder using electric blender. To prepare the plant extracts, 1000 g of the powdered leaves were dissolved in 1000 ml of distilled water. The mixture was thoroughly stirred overnight and then filtered through a double-layered muslin cloth to create a stock solution [23]. The stock solution was then dried using a rotary evaporator (Generic RE-201D rotary evaporator) and prepared in two concentrations (10,000 ppm and 20,000 ppm) for

testing against the isolated post-harvest fungi. These plant extracts, known for their effectiveness against fungal pathogens, were evaluated *In vitro* using the Poisoned Food Technique [24] and Potato Dextrose Agar (PDA) as the base medium.

Three replicates were maintained for both the test pathogens and the control (without plant extract addition). The petri plates were inverted and incubated at 28 ± 2°C and 65% to 70% RH. Observations on radial mycelial growth and percent inhibition of the test fungi were recorded at 24-hour intervals until the test pathogen on the untreated control plate completely covered the medium [25].

The percentage of inhibition for the test pathogen was determined as described by [26].

$$\text{Percent inhibition} = \left(\frac{C - T}{C} \right) \times 100$$

Where,

C = Growth of the test fungus in untreated control plates

T = Growth of the test fungus in treated plates

Essential oils of *Ocimum tenuiflorum*, *Eucalyptus globulus*, *Azadirachta indica*, *Pongamia pinnata*, *Cymbopogon citratus*, *Syzygium aromaticum*, *Mentha piperita* L., and *Punica granatum* L. were procured from local market and tested for the antifungal activity in similar way as performed for plant extracts. The concentration of essential oils used was 1500 and 2000 ppm. Percent inhibition were also calculated for the effect of essential oils.

2.4 Statistics

The mean, standard error of mean and analysis of variance with Tukey's HSD post hoc analysis was done using IBM SPSS ver 24.

3. RESULTS AND DISCUSSION

Post-harvest diseases of banana fruits i.e. crown rot, anthracnose, cigar end rot, fungal scald, stem end rot, main stalk rot, and botryodiplodia finger rot are some of the major diseases on different banana cultivars in India [27]. These diseases cause significant losses during storage, transportation, and marketing, affecting different banana cultivars. Globally, postharvest diseases account for 10-30% of total crop yield losses, and

in some perishable crops, especially in developing countries, they can destroy over 30% of the yield [28].

3.1 Isolation and Identification of Post-harvest Fungi in Banana

Table 1 outlined various postharvest diseases affecting *Musa paradisiaca* (banana) and their respective causative agents along with their accession numbers. The isolation of pathogens from post-harvested *Musa paradisiaca* fruits revealed several significant diseases. *Fusarium napiforme* was identified as the pathogen responsible for Fusarium wilt, also known as Panama disease, which severely affected the banana plants. *Talaromyces atroseus* was found to cause fruitlet core rot, compromising the quality and marketability of the fruit. *Cladosporium cladosporioides* led to crown rot, a common post-harvest disease resulting in significant losses during storage and transport. Another *Fusarium* species, *Fusarium sp.*, was also identified as a cause of Fusarium wilt or Panama disease, similar to *Fusarium napiforme*. Additionally, *Fusarium equiseti* was found to contribute to Fusarium wilt or Panama disease, further highlighting the widespread impact of this disease on banana production. Similar results were obtained by Sarkar, Girisham [14] who also reported the incidence of *Cladosporium cladosporioides* and *Fusarium sp.* from post-harvest fruits of banana from the Warangal region. Aradhana Pal, Singh [29] also reported a total of 12 fungi isolated from diseased banana fruit samples viz. *Aspergillus flavus*, *Aspergillus niger*, *Alternaria spp.*, *Botryodiplodia theobromae*, *Colletotrichum musae*, *Fusarium equiseti*, *Fusarium moniliforme*, *Fusarium oxysporum*, *Fusarium solani*, *Mucor circinelloides*, *Penicillium spp.*, and *Rhizopus stolonifer* from Uttar Pradesh state. The species

of fungi infecting the fruit may vary depending on the cultivar of the crop, the host range of the fungi and geographical location as well as the environmental factors affecting the crop [30].

3.2 Bio Efficacy of Essential Oils and Plant Extracts on the Fungus

The bioefficacy of different plant extracts and essential oils at different concentrations was tested against the post-harvest pathogenic fungus of banana fruit. The essential oils and aqueous plant extracts significantly exhibited percent inhibition of the selected post-harvest pathogens of banana fruits at $P < 0.05$ level of significance. With the increase in the concentration of essential oils and plant extracts, the inhibition also increased. At 1500 ppm concentration, essential oils exhibited varying degrees of bio-efficacy against post-harvest fungal pathogens as depicted in Table 2. *Syzygium aromaticum* and *Mentha piperita* L. oils achieved 100% inhibition of *Fusarium equiseti*. *Punica granatum* L. oil was most effective against *Fusarium napiforme* (46.83%) and *Talaromyces atroseus* (73%). *Pongamia pinnata* oil showed high efficacy against *Talaromyces atroseus* (67%) and *Cladosporium cladosporioides* (52.31%).

At 2000 ppm concentration, clove and peppermint oils maintained 100% inhibition of *Fusarium equiseti* (Table 3). Pomegranate oil significantly inhibited *Cladosporium cladosporioides* (59.65%), while *Pongamia pinnata* oil effectively inhibited *Fusarium napiforme* (33.28%) and *Cladosporium cladosporioides* (50.47%). In the current study, the essential oils of *Ocimum tenuiflorum*, *Mentha piperita* and *Syzygium aromaticum* were effective in inhibiting the growth of the fungi.

Table 1. Isolated pathogens from the *Musa paradisiaca* post-harvested fruits

Sr No.	Name of Pathogen Isolated	Accession number	Disease name
1	<i>Fusarium napiforme</i>	OL711962	Fusarium wilt or Panama disease
2	<i>Talaromyces atroseus</i>	OL711963	Fruitlet core rot
3	<i>Cladosporium cladosporioides</i>	OL711965	Crown rot
4	<i>Fusarium sp.</i>	OL711966	Fusarium wilt or Panama disease
5	<i>Fusarium equiseti</i>	OL711967	Fusarium wilt or Panama disease

Table 2. Bio-Efficacy of the essential oils against post-harvested fungal pathogens at 1500 ppm oil concentrations

Percent inhibition	<i>Ocimum tenuiflorum</i>	<i>Eucalyptus citriodora hook</i>	<i>Azadirachta indica</i>	<i>Pongamia pinnata</i>	<i>Cymbopogon citratus</i>	<i>Syzygium aromaticum</i>	<i>Mentha piperita L.</i>	<i>Punica granatum L.</i>
<i>Fusarium napiforme</i>	24.25±0.1	11.71±0.12	19.74±0.11	21.74±0.1	16.23±0.11	41.81±0.08	26.76±0.1	46.83±0.07
<i>Talaromyces atroseus</i>	22±0.15	15±0.17	6±0.18	67±0.06	18±0.16	57±0.08	64±0.07	73±0.05
<i>Cladosporium cladosporioides</i>	35.13±0.04	1.53±0.06	5.64±0.06	52.31±0.03	49.74±0.03	42.05±0.04	55.9±0.03	30±0.04
<i>Fusarium sp.</i>	29.27±0.06	26.45±0.01	28.33±0.04	26.92±0.02	23.63±0.06	22.85±0.09	24.57±0.04	31.77±0.07
<i>Fusarium equiseti</i>	13±0.27	26±0.23	20±0.25	6±0.3	22±0.24	100±0a	100±0a	18±0.26

The readings are in Percentage inhibition. Any value followed by ± denotes standard error of the mean. The mean values with similar alphabets are similar means at p<0.05 level of significance

Table 3. Bio-Efficacy of the essential oils against post-harvested fungal pathogens at 2000 ppm oil concentrations

Percent inhibition	<i>Ocimum tenuiflorum</i>	<i>Eucalyptus citriodora hook</i>	<i>Azadirachta indica</i>	<i>Pongamia pinnata</i>	<i>Cymbopogon citratus</i>	<i>Syzygium aromaticum</i>	<i>Mentha piperita L.</i>	<i>Punica granatum L.</i>
<i>Fusarium napiforme</i>	30.77 ± 0.09	27.26 ± 0.1	20.74 ± 0.11	33.28 ± 0.09	35.29 ± 0.09	50.34 ± 0.07	47.33 ± 0.07	28.27 ± 0.1
<i>Talaromyces atroseus</i>	11 ± 0.4cd	9 ± 0.41bc	9 ± 0.41bd	34 ± 0.3	-1 ± 0.46	64 ± 0.16	42 ± 0.26	52 ± 0.22
<i>Cladosporium cladosporioides</i>	44.87 ± 0.03	35.69 ± 0.04	59.65 ± 0.02	50.47 ± 0.03	2.27 ± 0.87	31.29 ± 0.04	40.48 ± 0.03	10.12 ± 0.05
<i>Fusarium sp.</i>	15.35 ± 0.31	41.73 ± 0.21bc	40.55 ± 0.22c	31.89 ± 0.25	36.61 ± 0.23a	37.8 ± 0.23a	42.52 ± 0.21b	48.03 ± 0.19
<i>Fusarium equiseti</i>	11.89 ± 0.6	24.32 ± 0.51b	18.67 ± 0.55	7.37 ± 0.63	23.19 ± 0.52b	100 ± 0a	100 ± 0a	1.73 ± 0.66

The readings are in Percentage inhibition. Any value followed by ± denotes standard error of the mean. The mean values with similar alphabets are similar means at p<0.05 level of significance

Table 4. Bio-Efficacy of the aqueous plant extracts against post-harvested fungal pathogens at 10000 ppm concentrations

Percent inhibition	<i>Ocimum tenuiflorum</i>	<i>Eucalyptus citriodora hook</i>	<i>Azadirachta indica</i>	<i>Pongamia pinnata</i>	<i>Cymbopogon citratus</i>	<i>Syzygium aromaticum</i>	<i>Mentha piperita L.</i>	<i>Punica granatum L.</i>
<i>Fusarium napiforme</i>	43.34 ± 0.15	35.57 ± 0.17a	63.34 ± 0.1	50.01 ± 0.13	44.45 ± 0.15	54.45 ± 0.12	65.56 ± 0.09	35.57 ± 0.17a
<i>Talaromyces atroseus</i>	72.13 ± 0.59ghi	56.52 ± 0.92cd	68.79 ± 0.66begi	65.44 ± 0.73aef	60.98 ± 0.83df	54.29 ± 0.97c	66.56 ± 0.71ab	73.24 ± 0.57gh
<i>Cladosporium cladosporioides</i>	55.41 ± 0.94begi	44.26 ± 1.18	64.33 ± 0.76h	57.64 ± 0.9adef	58.75 ± 0.87cfhi	28.65 ± 1.51	56.52 ± 0.92abc	52.06 ± 1.02dg
<i>Fusarium sp.</i>	73.43 ± 0.04	54.61 ± 0.07	65.68 ± 0.05	84.5 ± 0.02	57.93 ± 0.06	67.9 ± 0.05	77.86 ± 0.03	60.15 ± 0.06
<i>Fusarium equiseti</i>	13.65 ± 0.13	14.76 ± 0.13	11.44 ± 0.13	16.97 ± 0.13ade	16.97 ± 0.13cef	16.97 ± 0.13bdf	16.97 ± 0.13abc	12.55 ± 0.13

The readings are in Percentage inhibition. Any value followed by ± denotes standard error of the mean. The mean values with similar alphabets are similar means at p<0.05 level of significance

Table 5. Bio-Efficacy of the aqueous plant RNA extracts against post-harvested fungal pathogens at 20000 ppm concentrations

Percent inhibition	<i>Ocimum tenuiflorum</i>	<i>Eucalyptus citriodora hook</i>	<i>Azadirachta indica</i>	<i>Pongamia pinnata</i>	<i>Cymbopogon citratus</i>	<i>Syzygium aromaticum</i>	<i>Mentha piperita L.</i>	<i>Punica granatum L.</i>
<i>Fusarium napiforme</i>	35.57 ± 0.17	31.12 ± 0.18	56.67 ± 0.11	44.45 ± 0.15	34.46 ± 0.17	45.56 ± 0.14	67.78 ± 0.08	32.23 ± 0.18
<i>Talaromyces atroseus</i>	72.13 ± 0.59di	62.1 ± 0.8befg	65.44 ± 0.73bfhkm	68.79 ± 0.66ahij	66.56 ± 0.71cgjlm	53.18 ± 0.99	69.9 ± 0.64abcd	63.21 ± 0.78cekl
<i>Cladosporium cladosporioides</i>	72.13 ± 0.59cfg	57.64 ± 0.9	75.47 ± 0.52bef	68.79 ± 0.66cd	67.67 ± 0.68d	73.24 ± 0.57eg	77.7 ± 0.47ab	79.93 ± 0.43a
<i>Fusarium sp.</i>	66.92 ± 0.14a	50.38 ± 0.21	61.4 ± 0.16	77.94 ± 0.09	54.79 ± 0.19	66.92 ± 0.14a	72.43 ± 0.12	55.89 ± 0.19
<i>Fusarium equiseti</i>	15.87 ± 0.13cef	15.87 ± 0.13adeg	13.65 ± 0.13	15.87 ± 0.13bdf	13.65 ± 0.13g	9.23 ± 0.14	15.87 ± 0.13abc	11.44 ± 0.13

The readings are in Percentage inhibition. Any value followed by ± denotes standard error of the mean. The mean values with similar alphabets are similar means at p<0.05 level of significance.




Fig. 1. Phylogenetic analysis of *Fusarium napiforme* registered as accession OL711962




Fig. 2. Phylogenetic analysis of *Talaromyces atroseus* registered as accession OL711963




Fig. 3. Phylogenetic analysis of *Cladosporium cladosporioides* registered as accession OL711965




Fig. 4. Phylogenetic analysis of *Fusarium* sp. registered as OL711966




Fig. 5. Phylogenetic analysis of *Fusarium equiseti* registered as OL711967

In a similar bioassay, *Ocimum basilicum* oil demonstrated fungistatic effects on the mycelial growth of *Fusarium moniliforme*, *Botrydiploidiatheobromae*, and *Colletotrichum sp.* at a low concentration of 1.5 ml/L (0.15% v/v) [31]. Researchers in Sri Lanka identified *Lasiodiplodia theobromae*, *Fusarium proliferatum*, and *Colletotrichum musae* as crown rot pathogens in bananas, which could act alone or in combination, with greater severity when combined. *Cymbopogon nardus* and *Ocimum basilicum* oils effectively inhibited *Colletotrichum musae* and *Fusarium proliferatum* at 0.2–0.6% concentrations, with even lower doses effective in liquid bioassays. These oils proven synergistic effects in *in-vivo* tests [32]. They reported ultrastructural changes, including plasmalemma disruption, mitochondrial deformation, and alterations in hyphal wall thickness, were observed after treatment, indicating the mode of action of *Cymbopogon nardus* essential oil. Similar results were obtained by Singh and Tripathi [17] with essential oils from *Cinnamomum zeylanicum*, *Azadirachta indica*, and *Mentha arvensis* exhibit 100% efficacy against postharvest banana pathogens. *Cinnamomum zeylanicum* oil was fungistatic at 100 ppm and fungicidal at 200 ppm.

At 10000 ppm concentration, aqueous plant extracts demonstrated significant efficacy against post-harvest fungal pathogens (Table 4). *Azadirachta indica* extract showed the highest inhibition of *Fusarium napiforme* (63.34%) and *Talaromyces atrovirens* (68.79%). *Mentha piperita* L. and *Punica granatum* L. extracts notably inhibited *Fusarium sp.* (77.86% and 60.15% respectively). At 20000 ppm concentration, neem extract continued to show high efficacy against *Fusarium napiforme* (56.67%) and *Cladosporium cladosporioides* (75.47%) (Table 5). Peppermint extract effectively inhibited *Fusarium napiforme* (67.78%) and *Talaromyces atrovirens* (69.9%), while pomegranate extract showed notable inhibition of *Cladosporium cladosporioides* (79.93%).

Similar results were obtained by Pant, Manandhar [33] using three botanicals (*Azadirachta indica*, *Justicia adhatoda*, and *Eucalyptus globules*) tested against *Fusarium oxysporum*. *Eucalyptus* was the most effective (36.67% inhibition at 10%), followed by *Justicia adhatoda* (9.10%) and *Neem* (5.32%). Another study demonstrated the antifungal efficacy of *Ocimum gratissimum* and *Moringa oleifera* leaf extracts against eleven different

fungi, including *Penicillium digitatum*, *Saccharomyces cerevisiae*, *Aspergillus niger*, and *Fusarium oxysporum* among others. The leaf extracts of *O. gratissimum* and *M. oleifera* demonstrated inhibitory effects on *Aspergillus* and *Fusarium* species, suggesting their potential as safe and effective alternatives to chemical fungicides for controlling post-harvest fruit deterioration [34].

Biological control of post-harvest diseases of bananas with natural compounds, antioxidants, inorganic salts, fungicides, or even biosurfactants have already been studied. The combination of such compounds along with essential oils and plant extracts could also enable the improvement of the biological control. Once the inhibition activity has been confirmed under controlled conditions, it is crucial to conduct tests under natural infestation conditions and through real export scenarios to evaluate the true effectiveness of the biocontrol strategy. The current experiments were performed under rigorous conditions, using artificial inoculations using the high inoculum levels of the pathogenic species to study the effectiveness against them. These severe conditions are rarely encountered in practical situations, suggesting that the protection level may be even higher under natural infestation.

4. CONCLUSION

Based on our findings, essential oils proved more effective than plant extracts in controlling the pathogenic fungi isolated from the specified niche. Given the significant degree of infection by these fungi, combining essential oils with other compounds such as metal nanoparticles or biosurfactants could further enhance their efficacy in combating post-harvest diseases. While our experiments were confined to laboratory tests, it is imperative to conduct field trials for real-world validation. These results offer valuable insights for implementing sustainable strategies to control post-harvest fungi.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Maseko KH, Regnier T, Meiring B, Wokadala OC, Anyasi TA. Musa species variation, production, and the application of its processed flour: A review. *Scientia Horticulturae*. 2024;325:112688.
2. Sathiya R, Naveenkumar M, Senthilnathan S, Grunathan S, Devi MN, Banumathy V. Growth and instability in area, production and productivity of Banana in Tamil Nadu. *International Journal of Plant & Soil Science*. 2022;34(20):592-8.
3. Canton H. Food and agriculture organization of the United Nations—FAO. The Europa directory of international organizations 2021: Routledge. 2021;297-305.
4. OECD/FAO. *Agricultural Outlook 2023-2032*; 2023.
5. Falana OB, Durodola OI. Multimodal remote sensing and machine learning for precision agriculture: A review. *Journal of Engineering Research and Reports*. 2022;23(8):30-34. Available: <https://doi.org/10.9734/jerr/2022/v23i8740>
6. Picon A, Alvarez-Gila A, Seitz M, Ortiz-Barredo A, Echazarra J, Johannes A. Deep convolutional neural networks for mobile capture device-based crop disease classification in the wild. *Computers and Electronics in Agriculture*. 2019;161:280-90.
7. Ghorbani R, Wilcockson S, Koocheki A, Leifert C. Soil management for sustainable crop disease control: A review. *Environmental Chemistry Letters*. 2008;6:149-62
8. Viljoen A, Ma L-J, Molina AB. CHAPTER 8: Fusarium wilt (panama disease) and monoculture in banana production: Resurgence of a century-old disease. *Emerging plant diseases and global food security: Am Phytopath Society*. 2020;159-84.
9. Thangavelu R, Mostert D, Gopi M, Devi PG, Padmanaban B, Molina A, et al. First detection of Fusarium oxysporum f. sp. cubense tropical race 4 (TR4) on Cavendish banana in India. *European Journal of Plant Pathology*. 2019;154:777-86.
10. Thangavelu R, Mustafa M. First report on the occurrence of a virulent strain of Fusarium wilt pathogen (Race-1) infecting Cavendish (AAA) group of bananas in India. *Plant Disease*. 2010;94(11):1379-.
11. Godana EA, Yang Q, Zhang X, Zhao L, Wang K, Dhanasekaran S, et al. Biotechnological and biocontrol approaches for mitigating postharvest diseases caused by fungal pathogens and their mycotoxins in fruits: A review. *Journal of Agricultural and Food Chemistry*. 2023;71(46):17584-96.
12. Chakrabarti A, Ganapathi T, Mukherjee P, Bapat V. MSI-99, a magainin analogue, imparts enhanced disease resistance in transgenic tobacco and banana. *Planta*. 2003;216:587-96.
13. Jones D. *Diseases of Banana, Abacá and Enset*. (DR Jones ed.). CABI Publishing, Wallingford, UK, 544pp; 2000.
14. Sarkar S, Girisham S, Reddy S. Incidence of post-harvest fungal diseases of banana fruit in Warangal market. *Indian Phytopathology*. 2009;62(1):103-5.
15. Sarkar S, Girisham S, Reddy S. Some new postharvest fungal diseases of banana. *Annal Plant Sci*. 2013;2:149-52.
16. Snehalatharani A, Devappa V, Sangeetha C. Postharvest diseases of banana and their management. *Postharvest Handling and Diseases of Horticultural Produce: CRC Press*. 2021;201-10.
17. Singh R, Tripathi P. Cinnamomum zeylanicum essential oil in the management of anthracnose of banana fruits. *JIPBS*. 2015;2(3):290-9.
18. Abd-Alla M, El-Gamal N, El-Mougy N, Abdel-Kader M. Post-harvest treatments for controlling crown rot disease of Williams banana fruits (*Musa acuminata* L.) in Egypt. *Plant Pathology and Quarantine*. 2014;4(1):1-12.
19. Nicosia MGLD, Pangallo S, Raphael G, Romeo FV, Strano MC, Rapisarda P, et al. Control of postharvest fungal rots on citrus fruit and sweet cherries using a pomegranate peel extract. *Postharvest Biology and Technology*. 2016;114:54-61.
20. Leyronas C, Duffaud M, Nicot PC. Compared efficiency of the isolation methods for Botrytis cinerea. *Mycology*. 2012;3(4):221-5.
21. Saitoh K-i, Togashi K, Arie T, Teraoka T. A simple method for a mini-preparation of fungal DNA. *Journal of General Plant Pathology*. 2006;72:348-50.
22. Tamura K, Stecher G, Kumar S. MEGA11: molecular evolutionary genetics analysis

- version 11. Molecular biology and evolution. 2021;38(7):3022-7.
23. Sahi S, Habib A, Ghazanfar M, Badar A. In vitro evaluation of different fungicides and plant extracts against *Botryodiplodia theobromae*, the causal agent of quick decline of mango. *Pakistan Journal of Phytopathology*. 2012;24(2):137-42.
 24. Nene YL, Thapliyal PN. Fungicides in plant disease control: International Science Publisher; 1993.
 25. Jagtap G, Dhavale M, Dey U. Evaluation of natural plant extracts, antagonists and fungicides in controlling root rot, collar rot, fruit (brown) rot and gummosis of citrus caused by *Phytophthora* spp. in vitro. *Sci J Microbiol*. 2012;1(2):27-47.
 26. Vincent J. Distortion of fungal hyphae in the presence of certain inhibitors. *Nature*. 1947;159(4051):850-.
 27. Patil P, Devi SP, Gutam S, Uma S, Kumar PS, Ladaniya M, et al. Annual Report-2020 of ICAR-All India Coordinated Research Project on Fruits, ICAR-Indian Institute of Horticultural Research, Bengaluru; 2020.
 28. Agrios GN. *Plant pathology*: Elsevier; 2005.
 29. Aradhana Pal AP, Singh R, Pooja Singh PS, Tripathi N. Some studies on post-harvest pathogens of banana from Gorakhpur (UP); 2017.
 30. Drenth A, Guest DI. Fungal and oomycete diseases of tropical tree fruit crops. *Annual Review of Phytopathology*. 2016;54: 373-95.
 31. Dube S, Upadhyay P, Tripathi S. Antifungal, physicochemical, and insect-repelling activity of the essential oil of *Ocimum basilicum*. *Canadian journal of botany*. 1989;67(7):2085-7.
 32. Anthony S, Abeywickrama K, Dayananda R, Wijeratnam S, Arambewela L. Fungal pathogens associated with banana fruit in Sri Lanka, and their treatment with essential oils. *Mycopathologia*. 2004;157:91-7.
 33. Pant B, Manandhar S, Manandhar C, Baidhya S. In vitro evaluation of fungicides and botanicals against *Fusarium oxysporum* f. sp. *cubense* of banana; 2020.
 34. Kinge TR, Besong PN. Fungi diversity on some fruits and biological control using two plants extracts. *Journal of Advances in Biology & Biotechnology*. 2021;24(4):24-38.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/119963>