



Ethnopharmacological Insights into the Use of *Chenopodium ambrosioides* for Cough Treatment among Togolese Traditional Healers

Kodjo Akpadja ^{a,b*}, Koffi Koudouvo ^{a,b},
Aboudou Azizou Dahounom ^{a,b}, Komlanvi Esseh ^{a,b},
Hafez Ouro-Djéri ^{a,b}, Kpatcha Edjarew Poro ^c,
Komlantsè Togbenou ^{a,b}, Tchadjobo Tchacondo ^c
and Amégnona Agbonon ^{a,b}

^a Center for Research and Training on Medicinal Plants (CERFOPLAM, Faculty of Sciences, University of Lomé, 01BP 1515 Lomé 01, Togo.

^b Laboratory of Physiology and Pharmacology, Faculty of Sciences, University of Lomé, 01BP, 1515 Lomé 01, Togo.

^c Laboratory of Biomedical, Food and Environmental Health Sciences (LaSBASE), ESTBA, University of Lomé, 01BP 1515 Lomé 01, Togo.

Authors' contributions

This work was carried out in collaboration among all authors. Authors KA and KK designed the study. Authors KA and AAD participated in data collection. Author KA compiled the information, analysed the data and drafted the initial manuscript. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ejmp/2024/v35i61217>

Open Peer Review History:
This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124644>

Original Research Article

Received: 15/08/2024

Accepted: 08/10/2024

Published: 13/10/2024

*Corresponding author: E-mail: akpadjakodjo2018@gmail.com;

Cite as: Akpadja, Kodjo, Koffi Koudouvo, Aboudou Azizou Dahounom, Komlanvi Esseh, Hafez Ouro-Djéri, Kpatcha Edjarew Poro, Komlantsè Togbenou, Tchadjobo Tchacondo, and Amégnona Agbonon. 2024. "Ethnopharmacological Insights into the Use of *Chenopodium Ambrosioides* for Cough Treatment Among Togolese Traditional Healers". *European Journal of Medicinal Plants* 35 (6):172-86. <https://doi.org/10.9734/ejmp/2024/v35i61217>.

ABSTRACT

Togo has an ethnobotanical potential with a great diversity of medicinal plants widely used to treat human diseases. Cough associated with respiratory diseases is one of the pathologies traditionally treated with medicinal plants in Togo. *Chenopodium ambrosioides* (L.) is a species used in the traditional treatment of cough in Togo. The aim of this study was to document the ethnopharmacological uses of *Chenopodium ambrosioides* in Togolese traditional medicine for treatment of cough. An ethnopharmacological survey was conducted using the semi-structured interview method among Traditional Medicine Practitioners (TMPs) from six health districts spread across four health regions. The ethnopharmacological and ethnobotanical data collected were analyzed using the User Value (UV) and Frequency of Citation (FC) of medicinal plants associated with the species. A total of 127 TMPs participated in this study and 96 (76%) of them were male and 31 (34%) were female. 73% of the informants claimed usually use *Chenopodium ambrosioides* in the cough treatment compared to 27% who claimed to not use it to treat cough. *Chenopodium ambrosioides* was used in combination with fifty-two (52) other botanical species in the treatment of cough and in the study area, *Abrus precatorius* Linn. (13.59%) followed by *Ocimum gratissimum* Linn. (11.58%), *Zingiber officinale* Rosc. Trans. Linn. (9.73%), *Mangifera indica* Linn. (6.21%), *Eucalyptus citriodora* Hooker (4.87%), *Xylopi aethiopica* (Dunal) A. Rich. (4.36%) and *Eugenia caryophyllata* Thunb. (4.19%) were the most popular plants used for cough treatment among the plants recorded. Sixteen (16) other pathologies including pulmonary infections are also treated by *Chenopodium ambrosioides* (L.). Abscesses, intestinal parasitic diseases, incurable wounds, stomach aches and headaches are the most pathologies commonly treated by this plant. The whole plant (72%) and leaves (22%) are the primary parts utilized for medicinal purposes. Decoction (77.50%) is the main preparation method, with oral (80.83%) and cutaneous (10.83%) routes being the most common modes of administration. The findings from this study provide a solid ethnopharmacological foundation for future pharmacological and toxicological investigations into *Chenopodium ambrosioides* (L.).

Keywords: *Chenopodium ambrosioides*; cough; ethnopharmacology; Togo.

1. INTRODUCTION

“Cough is a protective immune response of the respiratory airways to expel or eliminate irritants and mucous. It is the most common symptom for which individuals seek medical advice” [1,2]. “Cough is the commonest symptom leading patients to consult with their doctor” [3]. “The epidemiology of cough can be neatly divided into these two diagnostic subgroups: acute cough, which is usually due to viral respiratory tract infection, and chronic cough, which may be arbitrarily defined as cough lasting longer than eight weeks” [3,4]. “The common causes of cough may be either of viral, or bacterial or chemical origin that include common cold, chronic bronchitis, rhinitis, sinusitis, asthma, environmental agents smoke, dust, angiotensin converting enzymes inhibitors and synthetic opioid analgesic” [5]. “Most coughs are acute and self-limiting, secondary to respiratory tract infection, but when cough becomes chronic (8 weeks duration) it can account for up to one third of all referrals to chest physicians” [6]. “Chronic cough has a major effect on the quality of life, and patients consume substantial

healthcare resources, often seeing several specialists over many years. It may prove diagnostically difficult, requiring multiple investigations and treatment trials, with diagnostic success quoted as ranging from 100% in a general respiratory clinic setting to 58% in the specialist cough clinic setting” [6,7]. “According to World Health Organization 80% of the Asian and African population relies on the medicinal plants for prevention and treatment of various diseases including cough” [8,9]. “Different communities and ethnic groups use medicinal plants, animal and minerals to treat different diseases. The people of the rural areas used traditional remedies to treat diseases” [2]. “They use traditional medicines due to i) knowledge from parents with firm belief, ii) knowledge of usage without consulting a physician, and iii) general perception of safety and economy” [2]. “*Chenopodium ambrosioides*, commonly called Mexican tea, is an aromatic plant belonging to Chenopodiaceae family and is known for its virtues” [10]. “Some research works have demonstrated that this medicinal plant is widely used by Togolese communities to

treat several diseases including the treatment of abscesses, epilepsy, nausea, stomachache, skin infections” [10,11]. “Some studies showed that *Chenopodium ambrosioides* is used in the cough and respiratory infections management in several Africa countries such as Benin, Ghana, Gabon, Ivory coast, Maroc, Nigeria and Senegal” [12–17]. But few studies have documented the endogenous uses of the species in the treatment of respiratory diseases in Togo. According to our literature review, no study on the endogenous knowledge of the uses of Togolese plants and *Chenopodium ambrosioides* in the treatment of cough in Togo is available. This is the main motivation of this study with the aim of documenting the uses of *Chenopodium ambrosioides* in the treatment of different forms of cough by traditional medicine practitioners in Togo. The objective of the study was to document and quantitatively analysed the traditional cough management recipes based on *Chenopodium ambrosioides* and preserve the traditional knowledge of the Togolese community for later uses. This ethnopharmacological study of *Chenopodium ambrosioides* is reported in Togo for the first time to treat cough and respiratory infections.

2. METHODOLOGY OF THE STUDY

2.1 Study Area

Togo is a Western African country surrounded in the North by the Republic of Burkina Faso, in the East by the Republic of Benin, in the West by the Republic of Ghana and in the South by the Atlantic Ocean. Togo has been divided into five economic regions from North to South: Savannah region, Kara region, Central region, Plateaux region, and Maritime region. After the general survey of the population in 2022 [18], Togo has an estimated population of 8 095 498 residents including 4 150 988 females (51,3%) with a rate of population growth of 2,3%. Our study area has an estimated population of 2 270 209 residents [18]. According to the organization of the health ministry, Togo has thirty-nine health districts grouped into six health regions such as Savannah, Kara, Central, Plateaux, Maritime and Grand Lome [19]. The present ethnopharmacological study was carried out in six health districts including Golfe, Ave, Yoto, Vo, Ogou and Tchamba (Fig. 1). Many languages are spoken in our study area but the most popular are Ewe, Mina, Ife and Tchamba.

2.2 Data Collection Period and Data Collection Forms

The survey was carried out from August 2021 to July 2022, to identify and record ethnopharmacological treatment of cough by traditional healers using *Chenopodium ambrosioides* alone or with other medicinal plants. The study objectives were explained to the healers and the most frequently asked questions were related to the local name of *Chenopodium ambrosioides*, medicinal uses in the cough management, parts used of the plant, the method of preparation, complete ingredients of the recipes, administration route and adverse effects. The data collection forms were prepared and explained to the informants in their local language before the interview [20–23].

2.3 Interview Techniques with the Traditional Healers

The data was collected through individual interviews following the semi-structured form previously prepared. The approach was in a first time, the group meeting with all the healers of the health district before the individual interview survey [21,24,25]. Conversations with the healers were used to obtain information on the uses of *Chenopodium ambrosioides* in the management of cough and respiratory infections. The healers that consented were asked to give their knowledge about the respiratory diseases they used *Chenopodium ambrosioides* against, the method of preparation of the remedy, details of administration, including the approximate amounts and number of doses per day or week, the composition of the remedy, the adverse effects of the remedy and how to treat these adverse effects. The conversations were built on trust with the common goal to improve the health situation in Togo and to preserve and increase the knowledge on medicinal plants, especially on *Chenopodium ambrosioides*.

2.4 Data Processing and Data Analysis

Analytical Flora of Benin [26] and the online database of PROTA4U (www.prota4u.org), were used for the botanical identification of species of plants species collected. An ethnobotanical and ethnopharmacological studies were carried out following the studies of [20,24,27,28]. The local names confirmation was carried out with reference to [26,29], West Africa Monograph Dictionary [30] and to the West African Pharmacopoeia [31]. Data collected was




Fig. 1. Map of Togo with focus on the survey areas

tabulates and Excel 2016 spreadsheet software were used to make simple calculations. The collected data from the survey questionnaire were analyzed using a quantitative method

including Use Value (UV) and Frequency of Citation (FC). This method enables us to make a consensus of the plant species which is frequently used to treat cough. FL indicates the

percentage of informants who claimed to use plant species for the same purpose regarding diseases' treatment. $FL (\%) = (N_p/N) \times 100$, where N_p indicates the number of the interviewed healers who claimed to use a plant species to treat a particular disease and N indicates the interviewed healers who used herbs as a medicine to cure any disease [24,32]. $FC (\%) = (\text{number of remedies containing the plant}/\text{total of remedies collected}) \times 100$. The Use Value, a quantitative method that demonstrates the relative importance of species known locally, was also calculated as follows: $UV = \sum U/n$, where UV is the Use value of a species, U is the number of citations per species, and n is the number of informants [25].

3. RESULTS AND DISCUSSION

A total of 127 participants was interviewed on average of 10 minutes. All the products of natural origin associated with *Chenopodium ambrosioides* were considered as part of the traditional recipes that were used to treat cough and respiratory diseases.

3.1 Characteristics of Traditional Healers

Of the 127 traditional healers surveyed, 96 were men (76%) and 31 were women (24%). Most of

them (89%) inherited the profession of "traditional healer" from their parents. This finding is in agreement with that of Dahounom et al., in Togo [33] and Umair et al., in Pakistan [34]. Sixty-two percent (62%) of the respondents claimed to have been practicing for 11–50 years (Table 1) with the majority (89%) having acquired the knowledge through ancestral inheritance. These findings are similar to those of Orch et al., in Morocco [35] and Odeunmi et al., in Nigeria [36]. This profession requires a long experience for the practitioner to be able to identify plants for effective management of diseases. Traditional religion (53%) and Christianity (33%) were the predominant religions of the respondents. About 37% of the healers were in the age between 18 and 40 years, 54% between 41 and 70 years, and 9% between 71 and 105 years (Table 1). This age distribution confirms the findings in Togo [37] and is similar to the findings in Cameroon [38] where a significant dominance of elder or very aged healers was documented. It suggests that the profession of traditional healers was mainly done by elders. About healers' education level, 15% held a university degree, 38% attended high and middle school, 30% attended primary school, and 17% were illiterate (Table 1). This finding is in agreement with that of Odeunmi et al., in Nigeria [36].

Table 1. Sociodemographic data of the respondents

Parameters	Respondents N (%)
Gender	
Male	96 (76%)
Female	31 (24%)
Age range (Years)	
[18-40]	47 (37%)
[41-70]	69 (54%)
71 and above	11 (9%)
Educational level	
Illiterate	22 (17%)
Primary	38 (30%)
Secondary	48 (38%)
University	19 (15%)
Ethnic group	
Ewe	81 (64%)
Ifè	18 (14%)
Tchamba	13 (10%)
Kabyè	05 (4%)
Others*	10 (8%)
Religion group	
Endogenous religion	67 (53%)
Christianism	42 (33%)
Muslim	17 (13%)
Buddism	01 (1%)

Parameters	Respondents N (%)
Origin of the knowledge	
Familial inheritance	113 (89%)
Initiation from traditional healers	12 (9%)
Divine revelation	02 (2%)
Period of practice (Years)	
[0-10]	44 (35%)
[11-50]	79 (62%)
51 and above	04 (3%)

*Kotocoli (2%), Fon (2%), Nawoudba (2%), Anyanga (1%) and Moba (1%)

3.2 *Chenopodium ambrosioides* and Cough Treatment in the Study Area

Cough being a symptom associated with several diseases, especially respiratory diseases, it remains the most common pathology in our communities. For the cough treatment, our communities mainly use medicinal plants including Chenopodiaceae genus. In our study, 73% of traditional healers claim used *Chenopodium ambrosioides* to treat cough (Fig. 2). Similarly, various authors have carried out the ethnobotanical studies in different countries and they reported *Chenopodium ambrosioides* for cough management [17, 38–43]. Cough is a global issue so various traditional remedies have been reported by different communities. Some of the plants like *Abrus precatorius* Linn.; *Ocimum gratissimum* Linn.; *Zingiber officinale* Rosc. Trans. Linn.; *Mangifera indica* Linn.; *Eucalyptus citriodora* Hooker; *Xylopiya aethiopica* (Dunal) A. Rich. and *Eugenia caryophyllata* have also been reported in our study for the treatment of cough. This result is in agreement with the study carry out in Nigeria by Adesina et al., [40]. Similarly, various authors have carried out the ethnobotanical studies in different areas and they reported those species of plants for treating cough [17,36,43,44]. Our recent study

corroborated the reported cough suppressive effect of *Chenopodium ambrosioides* L. [17]. This study with traditional practitioners revealed a varied use of *Chenopodium ambrosioides*, often in the form of decoctions, infusions or inhalations, to relieve cough symptoms. These practices are rooted in ancestral traditions that transmit not only knowledge about medicinal plants, but also cultural and social practices that shape the perception and application of treatments.

3.3 Others Pathologies Treated by *Chenopodium ambrosioides* (L.) in the Study Area

In addition to its use to treat different forms of cough, participants in this study reported that *Chenopodium ambrosioides* L. is also used to treat 16 other pathologies including abscesses, intestinal parasitic diseases, incurable wounds, stomach-aches, headaches, dermatosis, angina and lung infections (Fig. 3). This result is in agreement with other studies carried out in Togo by Tettegah et al., and Ouadja et al., [11,25], in Nigeria by Adesina et al., [40], in Benin by Alitonou et al., [15], in Morocco by Orch et al., [35] then in South Korea by Lee et al., [45].




Fig. 2. *Chenopodium ambrosioides* and cough treatment in the study area




Fig. 3. Others pathologies treated by *Chenopodium ambrosioides* in the study area

3.4 Botanical Diversity and Ethnobotanical Indices of Plants Used in the Treatment of Cough Associated with Respiratory Diseases

Despite the relative high number of plants utilised for cough purpose, their ethnobotanical indices including FC (1-13%) and UV (0.03-0.64) were generally low (Table 2). In the study area, *Abrus precatorius* Linn. (13.59%) followed by *Ocimum gratissimum* Linn. (11.58%), *Zingiber officinale* Rosc. Trans. Linn. (9.73%), *Mangifera indica* Linn. (6.21%), *Eucalyptus citriodora* Hooker (4.87%), *Xylopia aethiopica* (Dunal) A. Rich. (4.36%) and *Eugenia caryophyllata* Thunb. (4.19%) were the most popular plants used for cough treatment among the plants recorded. Locally known as “dzindzinkoudze (ewe) or vivima (ewe) or bibignakou (Tchamba)”, *Abrus precatorius* (Linn.) has long been regarded as a potent remedy for relieving cough among the Togolese population [25,32,37]. Member of the genus *Abrus* are known to often be used for diverse ailments including respiratory diseases in Africa [43,46,47] and in Togo [37]. The alkaloidal constituents, which are often characteristically of the family Fabaceae including the genus *Abrus*, are known to significantly contribute to their diverse medicinal attributes [48]. In total 52 medicinal plant species belonging to 29 families were documented (Table 2). Caesalpiniaceae (5 species) and Euphorbiaceae (4 species) were the most dominant families, followed by Annonaceae, Asteraceae, Fabaceae, Rubiaceae and Rutaceae (3 species each) while other families were represented by one species only (Fig. 4). The utilization of medicinal plant species belonging to Caesalpiniaceae, Euphorbiaceae, Annonaceae, Asteraceae and Fabaceae families

was in agreement with ethnomedicinal flora reported from other parts of Togo and in other areas of the world [2,28,33,48–51]. This may be due to their wide distribution of plant species and their traditional uses known by the indigenous communities living in different parts of the world. Moreover, in a systematic review conducted by Lee et al., [45] in South Korea in 2023 on medicinal plants used to treat chronic cough, the most cited species (*Glycyrrhiza glabra*) with 61 citations is part of the Fabaceae family. This confirms the use of species from the main families listed in this study in the treatment of respiratory diseases including cough.

3.5 Plant Parts, Mode of Preparation and Rout of Administration

Even though all plant parts are significant in the treatment of different ailments; nevertheless, in the present study as shown in Fig. 5a, whole plant were the most commonly utilized part of *Chenopodium ambrosioides* with 72% application in traditional medicinal recipes, followed by leaves (22%) and others (root, seed, flower) with 5% application. Tettegah et al., 2009 and Ouadja et al., 2021 reported whole plants and leaves as commonly utilized plant part in herbal medicine used by the population of Togo [11,25]. However, it has been reported that the use of leaves is better for the survival of medicinal plants collected by herbalists compared to the collection of whole plant, roots and stem, which may cause severe threat to local flora [40].

To facilitate the treatment of cough and pathologies of respiratory system with *Chenopodium ambrosioides*, different preparation modes and administration are employed, namely decoction, Infusion, maceration, powder and others (fumigation).






Fig. 4. Families of medicinal plant species used to treat cough in the study area



a) Plant parts used

b) Mode of preparation



c) Rout of administration

Fig. 5. a) Plant parts; b) Mode of preparation; c) Rout of administration

Table 2. Medicinal plants used for the treatment of cough in Togo

Vernacular names	Species	Families	Citation	FC*(%)	UV*
Dzedzinkoudzen (ewe), Bibignakou (Tchamba)	<i>Abrus precatorius</i> Linn.	Fabaceae	81	13,59%	0,64
Dzogbeti (ewe)	<i>Ocimum gratissimum</i> Linn.	Lamiaceae	69	11,58%	0,54
Dotè (ewe)	<i>Zingiber officinale</i> Rosc. Trans. Linn.	Zingiberaceae	58	9,73%	0,46
Mangoti (ewe)	<i>Mangifera indica</i> Linn.	Anacardiaceae	37	6,21%	0,29
Ekalypti (ewe)	<i>Eucalyptus citriodora</i> Hooker	Myrtaceae	29	4,87%	0,23
Etso (ewe); Ekamanté (Tchamba)	<i>Xylopia aethiopica</i> (Dunal) A. Rich.	Annonaceae	26	4,36%	0,20
Plingota (ewe)	<i>Eugenia caryophyllata</i> Thunb.	Myrtaceae	25	4,19%	0,20
Zanguarati (ewe)	<i>Cassia siamea</i> Linn.	Caesalpiniaceae	19	3,19%	0,15
Zokon (mina)	<i>Indigofera pulchra</i> Willd.	Fabaceae	18	3,02%	0,14
Exeti (ewe)	<i>Zanthoxylum xanthoxyloides</i> Lam.	Rutaceae	17	2,85%	0,13
Nyimon (ewe)	<i>Sarcocephalus latifolius</i>	Rubiaceae	17	2,85%	0,13
Donti (ewe)	<i>Citrus sinensis</i>	Rutaceae	16	2,68%	0,13
Kpatima (ewe)	<i>Newbouldia laevis</i> Seem.	Bignoniaceae	14	2,35%	0,11
Tsaka-tsaka (ewe)	<i>Securinega virosa</i> (Reyb.) Baill.	Euphorbiaceae	13	2,18%	0,10
Yovovigbe (ewe)	<i>Moringa oleifera</i> Linn.	Moringaceae	11	1,85%	0,09
Ahamè (ewe)	<i>Ocimum canum</i> Sams	Lamiaceae	9	1,51%	0,07
Aloma (ewe)	<i>Vernonia colorata</i> (Willd.) Drake Bull.	Asteraceae	9	1,51%	0,07
Mahougen (ewe)	<i>Khaya senegalensis</i> (Desr.) A. Juss.	Meliaceae	9	1,51%	0,07
Avlokouigbe (ewe)	<i>Pleiocarpa pycnantha</i> (K. Sch.) Stapf.	Apocynaceae	8	1,34%	0,06
Ehlinvi (ewe)	<i>Phyllanthus amarus</i> Sch. et Th.	Euphorbiaceae	7	1,17%	0,06
Ewoti (ewe); Boudo (Tchamba)	<i>Parkia biglobosa</i> (Jacq.) Benth.	Mimosaceae	7	1,17%	0,06
Kiniti (ewe)	<i>Azadirachta indica</i> Linn.	Meliaceae	7	1,17%	0,06
Yovoninti (ewe)	<i>Cocos nucifera</i> Linn.	Arecaceae	7	1,17%	0,06
Akamati (ewe)	<i>Bridelia ferruginea</i> Benth.	Euphorbiaceae	6	1,01%	0,05
Damelatsu'nko (ewe); Efoulandikonkon (tchamba)	<i>Acanthospermum hispidum</i> (DC).	Asteraceae	6	1,01%	0,05
Pamplemousse	<i>Citrus Lemon</i>	Rutaceae	6	1,01%	0,05
Adako (ewe)	<i>Sorghum caudatum</i> Linn. var. colorans	Poales	5	0,84%	0,04
Anonsikan (ewe)	<i>Euphorbia hirta</i> Linn.	Euphorbiaceae	5	0,84%	0,04
Fefeti (ewe)	<i>Gardenia ternifolia</i> Sch. et Th.	Rubiaceae	5	0,84%	0,04
Hédjati (ewe)	<i>Harrisonia abyssinica</i>	Simaroubaceae	5	0,84%	0,04

Vernacular names	Species	Families	Citation	FC*(%)	UV*
Hehetsi (ewe)	<i>Anogeissus leiocarpus</i> Guill. et Pers.	Combretaceae	5	0,84%	0,04
Tsigbe (ewe)	<i>Cymbopogon citratus</i> Stapf.	Poaceae	5	0,84%	0,04
Ekloti (ewe)	<i>Piliostigma thonningii</i> (Sch.) Miln. Redh.	Caesalpiniaceae	4	0,67%	0,03
Fonti (ewe)	<i>Vitex doniana</i> Sweet	Verbenaceae	4	0,67%	0,03
Azingbe (ewe)	<i>Cassia rotundifolia</i> Linn.	Caesalpiniaceae	3	0,50%	0,02
Etritou (ewe), Forze (tchamba)	<i>Securidaca longepedunculata</i> Fres.	Polygalaceae	3	0,50%	0,02
Adikouti (ewe)	<i>Caesalpinia bonduc</i> Roxb.	Caesalpiniaceae	2	0,34%	0,02
Alangbati (ewe), Ditori (tchamba)	<i>Adansonia digitata</i> Linn.	Bombacaceae	2	0,34%	0,02
Avakofé (mina), Faux kinkeliba	<i>Cassia occidentalis</i> Linn.	Caesalpiniaceae	2	0,34%	0,02
Somboegbé (ewe)	<i>Cleome viscosa</i> Linn.	Cucurbitaceae	2	0,34%	0,02
Tèctki (ewe)	<i>Tectona grandis</i> Linn.	Verbenaceae	2	0,34%	0,02
Adoubati (ewe)	<i>Carica papaya</i> Linn.	Caricaceae	1	0,17%	0,01
Agbanlan (mina)	<i>Uvaria chamae</i> P. Beauv.	Annonaceae	1	0,17%	0,01
Agnigli (ewe) ; Choechoere (Tchamba)	<i>Annona senegalensis</i> Pers.	Annonaceae	1	0,17%	0,01
Assiviatoe (ewe)	<i>Cola millenii</i> K. Schum.	Sterculiaceae	1	0,17%	0,01
Gbolobavi (ewe)	<i>Calotropis procera</i> (Ait.) Ait.	Asclepiadaceae	1	0,17%	0,01
Gnakpekpe (ewe)	<i>Kigelia africana</i> (Lam.) Benth.	Bignoniaceae	1	0,17%	0,01
Nawa (Kotocoli)	<i>Vernonia guineensis</i>	Asteraceae	1	0,17%	0,01
Nyamidua (Ashanti)	<i>Alstonia boonei</i> De Wild.	Apocynaceae	1	0,17%	0,01
Papatahé (ewe)	<i>Gomphrena celosioides</i> Mart.	Amaranthaceae	1	0,17%	0,01
Sifafa (ewe)	<i>Pavetta corymbosa</i> (DC) F.N. Will.	Rubiaceae	1	0,17%	0,01
Vénaviwogbe (ewe)	<i>Uraria picta</i> (Jacq.) DC	Fabaceae	1	0,17%	0,01

*FC = Frequency; *UV = Use Value

The decoction remains the most used mode (77,50%), because some questioned people believe that this mode allows to collect the most active ingredients and to mitigate or cancel the toxicity of certain recipes. Then come others preparation modes, such as: infusion (14,17%), maceration (5,83%) and powder (2,50%) (Fig. 5b). Oral (80.83%) and cutaneous (10.83%) (Fig. 5c) are the main routes of administration. The similar results were reported by Orch et al. in Morocco [35]. Decoction and infusion constitute the essentials mode of preparation and use recommended in traditional therapy.

4. CONCLUSION

This ethnopharmacological survey on cough treatment with *Chenopodium ambrosioides* in the four health regions of Togo, including: Grand Lome, Maritime, Plateaux and Centrale, revealed that traditional healers had an endogenous understanding of cough associated with the respiratory diseases. The present study documented anticough medicinal plant species. As the emergence of resistance of microorganisms that can cause cough such as *Streptococcus pneumonia* and *Mycobacterium tuberculosis* to antibiotics is a deep concern, Togolese flora must be explored for new bioactive anticough drugs. The commonly recommended strategy in the discovery of drugs is to evaluate plant extracts *in vitro* (antimicrobial activity) and *in vivo* (antitussive activity) to confirm their therapeutic potential. Keeping this in mind, our ethnopharmacological survey is the first step before confirmation *in vitro* and *in vivo*. Some studies are in progress to validate the antioxidant, antimicrobial, and antitussive effects of *Chenopodium ambrosioides* associated with some species documented in this study.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENTS

The authors are thankful to the people of the area, who participate in the study voluntarily. We are also grateful to the traditional healers of the four regions who participated in this survey. Our special thanks to the coordinator of the different association of traditional healers in the area of the survey. We are also grateful to the staff members of the herbarium of the University of Lome.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Irwin RS, Madison JM, Fraire AE. The cough reflex and its relation to gastroesophageal reflux. *Am J Med.* 2000;108:73–8. Available: [https://doi.org/10.1016/S0002-9343\(99\)00341-1](https://doi.org/10.1016/S0002-9343(99)00341-1)
2. Shaheed Benazir Bhutto University Sheringal Dir Upper Khyber Pakhtun Khwa, Khan M, Ali M, Shaheed Benazir Bhutto University Sheringal Dir Upper Khyber Pakhtun Khwa, Ullah R, Shaheed Benazir Bhutto University Sheringal Dir Upper Khyber Pakhtun Khwa, et al. Ethnopharmacological treatment of Cough in Piran, Malakand, Pakistan. *Boletin Latinoam Caribe Plantas Med Aromat.* 2021;20:203–14. Available: <https://doi.org/10.37360/blacpma.21.20.2.16>
3. Morice AH. Epidemiology of Cough. *Pulm Pharmacol Ther.* 2002;15:253–9. Available: <https://doi.org/10.1006/pupt.2002.0352>
4. Song W-J, Morice AH, Kim M-H, Lee S-E, Jo E-J, Lee S-M, et al. Cough in the Elderly Population: Relationships with Multiple Comorbidity. *PLoS ONE.* 2013;8:e78081. Available: <https://doi.org/10.1371/journal.pone.0078081>.
5. Duricek M, Nosakova L, Zatko T, Pecova R, Hyrdel R, Banovcin P. Cough reflex sensitivity does not correlate with the esophageal sensitivity to acid in patients with gastroesophageal reflux disease. *Respir Physiol Neurobiol.* 2018;257:25–9.

- Available:<https://doi.org/10.1016/j.resp.2018.03.011>.
6. Clare Decalmer S, Webster D, Alice Kelsall A, McGuinness K, Arthur Woodcock A, Ann Smith J. Chronic cough: How do cough reflex sensitivity and subjective assessments correlate with objective cough counts during ambulatory monitoring? *Thorax*. 2007;62:329–34.
Available:<https://doi.org/10.1136/thx.2006.067413>.
 7. Kelsall A, Decalmer S, McGuinness K, Woodcock A, Smith JA. Sex differences and predictors of objective cough frequency in chronic cough. *Thorax*. 2009;64:393–8.
Available:<https://doi.org/10.1136/thx.2008.106237>.
 8. World Health Organization. WHO global report on traditional and complementary medicine 2019. Geneva: World Health Organization; 2019.
 9. Bodeker G, Kronenberg F. A Public Health Agenda for Traditional, Complementary, and Alternative Medicine. *Am J Public Health*. 2002;92:1582–91.
Available:<https://doi.org/10.2105/AJPH.92.10.1582>.
 10. Ouadja B, Katawa G, Toudji GA, Layland L, Gbekley EH, Ritter M, et al. Anti-inflammatory, antibacterial and antioxidant activities of *Chenopodium ambrosioides* L. (Chenopodiaceae) extracts. *J Appl Biosci*. 2021;162:16764–94.
Available:<https://doi.org/10.35759/JABs.162.7>.
 11. Tettegah M, Eklou-Kadegbeku K, Aklikokou A, Agbonon A, De Souza C, Gbeassor M. Infected wound healing and antimicrobial effects of *Chenopodium ambrosioides* and *Mitracarpus scaber*. *Int J Biol Chem Sci*. 2009;3.
Available:<https://doi.org/10.4314/ijbcs.v3i3.45329>.
 12. Okhale Samuel Ehiabhi. Phytochemical and proximate analyses and thin layer chromatography fingerprinting of the aerial part of *Chenopodium ambrosioides* Linn. (Chenopodiaceae). *J Med Plants Res*. 2012;6.
Available:<https://doi.org/10.5897/JMPR11.201>.
 13. El Hafian M, Benlandini N, Elyacoubi H, Zidane L, Rochdi A. Étude floristique et ethnobotanique des plantes médicinales utilisées au niveau de la préfecture d'Agadir-Ida-Outanane (Maroc). *J Appl Biosci*. 2014;81:7198.
Available:<https://doi.org/10.4314/jab.v81i1.8>.
 14. Sousa ZL, De Oliveira FF, Da Conceição AO, Silva LAM, Rossi MH, Santos JS, et al. Biological activities of extracts from *Chenopodium ambrosioides* Lineu and *Kielmeyera neglecta* Saddi. *Ann Clin Microbiol Antimicrob*. 2012;11:20.
Available:<https://doi.org/10.1186/1476-0711-11-20>.
 15. Alitonou GA, Sessou P, Tchobo FP, Avlessi F, Yehouenou B, Menut C, et al. Chemical composition and biological activities of essential oils of *Chenopodium ambrosioides* L. collected in two areas of Benin; 2012.
 16. Togola A, Diallo D, Dembélé S, Barsett H, Paulsen BS. Ethnopharmacological survey of different uses of seven medicinal plants from Mali, (West Africa) in the regions Doila, Kolokani and Siby. *J Ethnobiol Ethnomedicine*. 2005;1:7.
Available:<https://doi.org/10.1186/1746-4269-1-7>.
 17. Loufoua BAE, Bassoueka DJ, Nsonde Ntandou GF, Nzonzi J, Etou-Ossibi AW, Ouamba JM, et al. Étude ethnobotanique, pharmacologique et phytochimique de quelques plantes médicinales congolaises à potentialité antitussive. *Phytothérapie*. 2015;13:377–83.
Available:<https://doi.org/10.1007/s10298-015-1005-4>.
 18. RGPH5 – INSEED; 2023.
Available:<https://inseed.tg/category/rgph5/> (accessed June 11, 2024).
 19. DSINISI. Statistical yearbook of the Ministry of Health.pdf; 2021.
 20. Koudouvo K, Karou DS, Kokou K, Essien K, Aklikokou K, Glietho IA, et al. An ethnobotanical study of antimalarial plants in Togo Maritime Region. *J Ethnopharmacol*. 2011;134:183–90.
Available:<https://doi.org/10.1016/j.jep.2010.12.011>.
 21. Koudouvo K, Esseh K, Denou A, Aziati T, Ajavon C, Afanyibo Y-G, et al. Sixth article: Ethno-pharmacological study of antimalarial medicinal recipes from Togo for the formulation of a

- phytomedicine for the treatment of malaria. Line; 2016.
22. Holaly GE, Simplicie KD, Charlemagne G, Kodjovi A, Kokou A, Tchadjobo T, et al. Ethnobotanical study of plants used in the treatment of diabetes in traditional medicine in the Maritime region of Togo. *Pan Afr Med J.* 2015;20. Available:<https://doi.org/10.11604/pamj.2015.20.437.5660>.
 23. Karou SD, Tchacondo T, Djikpo Tchibozo MA, Abdoul-Rahaman S, Anani K, Koudouvo K, et al. Ethnobotanical study of medicinal plants used in the management of diabetes mellitus and hypertension in the Central Region of Togo. *Pharm Biol.* 2011;49:1286–97. Available:<https://doi.org/10.3109/13880209.2011.621959>.
 24. Yaovi-Gameli A, Koffi K, Komlavi E, Amegnona A, Koffi T, Messanvi G. An Ethnobotanical Survey of Medicinal Plants used in the Preparation of “Atikédi”: Local Alcoholic Beverages Commonly Consumed in Lomé Togo. *Eur Sci J ESJ.* 2018;14:1. Available:<https://doi.org/10.19044/esj.2018.v14n33p1>.
 25. Ouadja B, Katawa G, Toudji GA, Layland L, Gbekley EH, Ritter M, et al. Anti-inflammatory, antibacterial and antioxidant activities of *Chenopodium ambrosioides* L. (Chenopodiaceae) extracts. *J Appl Biosci.* 2021;162:16764–94. Available:<https://doi.org/10.35759/JABs.162.7>.
 26. Flora Analytical of Benin. by Akoegninou, A. et al (Eds.). | C. Arden (Bookseller) ABA n.d. Available:<https://www.abebooks.fr/Flore-Analytique-Benin-Akoegninou-A-Eds/31636583905/bd> (accessed June 15, 2024).
 27. Koudouvo K, Esseh K, Denou A, Aziati T, Ajavon C, Afanyibo Y-G, et al. Sixth article: Ethno-pharmacological study of antimalarial medicinal recipes from Togo for the formulation of a phytomedicine for the treatment of malaria. Line; 2016.
 28. Effoe S, Gbekley EH, Mélila M, Aban A, Tchacondo T, Osseyi E, et al. Ethnobotanical study of food plants used in traditional medicine in the Maritime region of Togo. *Int J Biol Chem Sci* 2020;14:2837–53. Available:<https://doi.org/10.4314/ijbcs.v14i8.15>.
 29. Joshua AÉ, Adjanohoun ÉJ. Contribution to ethnobotanical and floristic studies in the People's Republic of Benin / E. J. Adjanohoun [et al.]. Paris: Agency for Cultural and Technical Cooperation; 1989.
 30. African Pharmacopoeia. Multilingual dictionary and monographs of the medicinal potential of African plants (West Africa) 2 volumes - Raphaël Eklun-Natey, Annie Balet. n.d.
 31. OOAS. la-pharmacopee-des-plantes-medicinales-de-lafrique-de-louest first edition; 2013. Available:<https://www.wahooas.org/web-oaas/sites/default/files/publications/2185/la-pharmacopee-des-plantes-medicinales-de-lafrique-de-louestok.pdf> (accessed June 10, 2024).
 32. Koukoura KK, Salifou TS, Gbekley EH, Pissang P, Effoe S, Tchacondo T, et al. Ethnobotanical survey of medicinal plants used in the treatment of vaginal and intestinal infections in the maritime region of Togo; 2022.
 33. Dahounom AA, Koudouvo K, Akpadja K, Esseh K, Ouro-Djeri H, Agbonon A, et al. Ethnobotanical and ethnopharmacological surveys on the endogenous knowledge of traditional medicine practitioners in the maritime region of Togo on *Chamaecrista rotundifolia* (pers.) Greene: Ethnobotanical and ethnopharmacological surveys on the endogenous knowledge of traditional medicine practitioners in the maritime region of Togo on *Chamaecrista Rotundifolia* (pers.) Greene. *J Rech Sci L'Université Lomé.* 2023;25:49–67.
 34. Umair M, Altaf M, Abbasi AM. An ethnobotanical survey of indigenous medicinal plants in Hafizabad district, Punjab-Pakistan. *PLoS ONE.* 2017;12:e0177912. Available:<https://doi.org/10.1371/journal.pone.0177912>.
 35. Orch H, Zidane L, Douira A. Ethnobotanical study of plants used in the treatment of respiratory diseases in a population bordering the forest of Izarene. *J Pharm Pharmacogn Res.* 2020;8:392–409.

- Available:https://doi.org/10.56499/jppres20.819_8.5.392.
36. Odebunmi CA, Adetunji TL, Adetunji AE, Olatunde A, Oluwole OE, Adewale IA, et al. Ethnobotanical survey of medicinal plants used in the treatment of COVID-19 and Related Respiratory Infections in Ogbomosho South and North Local Government Areas, Oyo State, Nigeria. *Plants*. 2022;11:2667. Available:<https://doi.org/10.3390/plants11192667>.
 37. Gbekley HE, Katawa G, Karou SD, Anani Sk, Tchadjobo T, Ameyapoh Y, et al. Ethnobotanical study of plants used to treat asthma in the Maritime Region in Togo. *Afr J Tradit Complement Altern Med AJTCAM*. 2017;14:196–212. Available:<https://doi.org/10.21010/ajtcam.v14i1.22>.
 38. Ladoh-Yemeda CF, Vandj T, Dibong SD, Mpondo EM, Wansi JD, Betti JL, et al. Étude ethnobotanique des plantes médicinales commercialisées dans les marchés de la ville de Douala, Cameroun. *J Appl Biosci*. 2016;99:9450–66. Available:<https://doi.org/10.4314/jab.v99i1.11>.
 39. Sbai-Jouilil H, Fadli A, Hafian ME, Ayad RE, Benharbit O, Zidane L. Floristic and Ethnobotanical Study of Medicinal Plants Used in the Treatment of Respiratory Diseases in Seksaoua Region (Western High Moroccan Atlas). *Annu Res Rev Bio*. 2017;17:1–10. Available:<https://doi.org/10.9734/ARRB/2017/36526>.
 40. Adesina S, Johnny I, Olayiwola G. Plants in Respiratory Disorders II- Antitussives, A Review. *Br J Pharm Res*. 2017;16:1–21. Available:<https://doi.org/10.9734/BJPR/2017/32974>.
 41. Mahboubi M. Management of acute cough by *Zataria multiflora* Boiss as an alternative treatment. *J Integr Med*. 2018;16:20–5. Available:<https://doi.org/10.1016/j.joim.2017.12.006>.
 42. Mathouet Ha, Aboughe Angone Sa, Mengome La, Eyele Mve Mba Ca, Rondi MLa, Souza Ab, Lamidi Ma. Ethnobotanical Study of Plants Used in Traditional Medicine for Respiratory Diseases in Gabon; 2014. Available:<https://doi.org/10.1016/j.pce.2022.103232>.
 43. Nyagumbo E, Pote W, Shopo B, Nyirenda T, Chagonda I, Mapaya RJ, et al. Medicinal plants used for the management of respiratory diseases in Zimbabwe: Review and perspectives potential management of COVID-19. *Phys Chem Earth Parts ABC*. 2022;128:103232. Available:<https://doi.org/10.1016/j.pce.2022.103232>.
 44. Kola-Mustapha AT, Yohanna KA, Ghazali YO, Ayotunde HT. Design, formulation and evaluation of *Chasmanthera dependens* Hochst and *Chenopodium ambrosioides* Linn based gel for its analgesic and anti-inflammatory activities. *Heliyon*. 2020;6:e04894. Available:<https://doi.org/10.1016/j.heliyon.2020.e04894>.
 45. Lee B, Kwon C-Y, Suh H-W, Kim YJ, Kim K-I, Lee B-J, et al. Herbal medicine for the treatment of chronic cough: A systematic review and meta-analysis. *Front Pharmacol*. 2023;14. Available:<https://doi.org/10.3389/fphar.2023.1230604>.
 46. Getachew S, Medhin G, Asres A, Abebe G, Ameni G. Traditional medicinal plants used in the treatment of tuberculosis in Ethiopia: A systematic review. *Heliyon*. 2022;8:e09478. Available:<https://doi.org/10.1016/j.heliyon.2022.e09478>.
 47. Gupta VK, Kaushik A, Chauhan DS, Ahirwar RK, Sharma S, Bisht D. Antimycobacterial activity of some medicinal plants used traditionally by tribes from Madhya Pradesh, India for treating tuberculosis related symptoms. *J Ethnopharmacol*. 2018;227:113–20. Available:<https://doi.org/10.1016/j.jep.2018.08.031>.
 48. Wang D, Wang S, Chen X, Xu X, Zhu J, Nie L, et al. Antitussive, expectorant and anti-inflammatory activities of four alkaloids isolated from Bulbus of *Fritillaria wabuensis*. *J Ethnopharmacol*. 2012;139:189–93. Available:<https://doi.org/10.1016/j.jep.2011.10.036>.
 49. Koudouvo K, Karou DS, Kokou K, Essien K, Aklidikou K, Glitho IA, et al. An ethnobotanical study of antimalarial plants

- in Togo Maritime Region. J Ethnopharmacol .2011;134:183–90.
Available:https://doi.org/10.1016/j.jep.2010.12.011.
50. Suroowan S, Mahomoodally MF. A comparative ethnopharmacological analysis of traditional medicine used against respiratory tract diseases in Mauritius. J Ethnopharmacol. 2016;177:61–80.
Available:https://doi.org/10.1016/j.jep.2015.11.029.
51. Aboudou AD, Koudouvo K. Ethnobotanical survey of medicinal plants used in the traditional management of infectious diseases in the Lomé-Commune Health Region of Togo. Eur Sci J ESJ. 2021;17:46–46.
Available:https://doi.org/10.19044/esj.2021.v17n21p46.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/124644>