



Carpal Tunnel Syndrome Caused by Palmaris Profundus Sharing Common Sheath with Median Nerve

**Nouha Baba^a, Senda Bellila^{a*}, Feten Sbei^a
and Mohamed Ali Sbai^a**

^a *Department of Plastic Surgery, University Hospital of Nabeul, Tunisia.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97318>

Case Report

Received: 07/01/2023
Accepted: 10/03/2023
Published: 14/03/2023

ABSTRACT

Carpal tunnel syndrome is the most common entrapment neuropathy in adults, usually caused by thickening of the flexor retinaculum. Occasionally, compression of the median nerve at the wrist may be caused by the presence of an intracanalicular tumor or anatomic variations. We report in this manuscript, the case of a carpal tunnel syndrome secondary to the presence of a palmaris profundus tendon included in a common sheath with the median nerve.

Keywords: *Carpal tunnel syndrome; median nerve; palmaris profundus.*

1. INTRODUCTION

Carpal tunnel syndrome is the most common entrapment neuropathy in adults, usually caused

by thickening of the flexor retinaculum, but it can also be secondary to various affections such as ganglion cysts, lipoma, synovial hypertrophy, bone fracture, vascular malformations or

*Corresponding author: E-mail: bellilasenda@gmail.com;

accessory muscles. Anatomic variations involving carpal tunnel and muscles in the forearm are well described by clinical and cadaveric studies [1,2]. Knowledge of the normal anatomy and anatomic variations is so important for any hand physician including orthopaedics and physiatrist for better understanding and managing these patients.

We describe in this article, an unusual case of the palmaris profundus found enclosed in a common fascial sheath with the median nerve, denoting an intimate relationship between the two structures, operated in the department of plastic surgery of University Hospital of Nabeul, Tunisia.

2. CASE PRESENTATION

A 48-year-old right-handed housewife, mother of 2 children, was referred to our consultation with a 3 year history of paresthesia in the territory of the median nerve of the right hand, nocturnal then diurnal as well as decrease of the grip strength. These symptoms were persistent despite symptomatic treatment (paracetamol and ibuprofene) prescribed by her family doctor.

Clinical examination revealed a median nerve with a grade 3 muscle strength on the thetheBMRC BMRC scale (British Medical Research Council) with no thenar atrophy and a grade 2 in the sensory scale. Tinel's sign was positive on percussion over the median nerve at the wrist with Phalen's test equally as positive.

Nerve conduction tests showed prolonged median motor and sensory latencies of distal origins, concluding to the diagnosis of a carpal tunnel syndrome.

The patient was operated under loco-regional anesthesia with a mini-incision approach. Upon surgery, release of the flexor retinaculum showed tendinous tissue that closely followed the nerve since before entrance of the carpal tunnel adhering intimately to it and seeming to have a common sheath with the nerve inside the tunnel, thus compressing it (Fig. 1).

We performed a dissection and release of the common sheath, though individualizing the nerve and the anomalous tendon (Fig. 2). The distal portion of the tendon was inserted into the palmar aponeurosis. We elected not to continue proximal dissection of the tendon in the forearm although we believe that it originated at a

considerable depth. Noted also was the absence of a palmaris longus tendon. The median nerve had an inflammatory aspect and was decreased in diameter, while the flexor tendon showed proliferative synovitis.



Fig. 1. Median nerve and palmaris profundus in the Common Sheath (CS). Flexor Retinaculum (FR) is released



Fig. 2. Individualization of Palmaris Profundus (PP) and Median Nerve (MN)

Postoperatively, pain and paresthesia disappeared and a sensory recovery graded at 4 was noted. No electromyogram was done as part of the follow-up.

3. DISCUSSION

Symptomatic compression on the median nerve occurs frequently and, most commonly, in the carpal tunnel. Anatomic variations in carpal tunnel are well documented in literature. However, there have been a few reports of median nerve compression caused directly by adherence of the tendon of palmaris profundus to the nerve. Some authors consider that it merely represents an anatomical variant of palmaris longus [1]. This muscle is of interest because of its variable anatomy and because of its inconsistency (absent in 17% of the general population) [3].

Palmaris profundus was first described in 1908 by Frohse and Frankel [4]. "Its frequency is about 0.18 % based on a study by Reimann et al. (1944) who found the muscle once in 530 limbs" [5]. It represents 17.8% of muscles involved with carpal tunnel syndrome [2]. "Palmaris profundus has been found with a proximally located muscle belly and also with a reversed, distally located belly" [6]. "This rare muscle is generally described as arising from the lateral edge of the radius (near the radial tuberosity), in its middle third, lateral to flexor digitorum superficialis and deep to pronator teres. Its tendon passes beneath the flexor retinaculum on the radial side of the median nerve, and after passing the carpal canal it broadens in the palm to insert into the deep surface of the palmar aponeurosis" [7, 8].

The similarity therefore of this muscle, found in our patient, to variable forms of palmaris longus is remarkable but they differed in one very important aspect from palmaris longus. The muscle is of special interest because it was found enclosed in a common fascial sheath with the median nerve. This unusual muscle in spite of its great resemblance to palmaris longus, may be better named "musculus comitans nervi mediani" [9], to denote its very important relationship to the median nerve, that of being its intimate traveling companion through the forearm and into the hand by way of the carpal canal.

4. CONCLUSION

Palmaris profundus muscle is a rare entity which could cause median nerve entrapment in carpal

tunnel, either by its tendon or its belly. In this case, standard surgical release of flexor retinaculum was sufficient to relieve symptoms. Other alternatives may include resection of the anomalous tendon.

CONSENT

Written consent of the patient has been collected and preserved by authors.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mitchell Ryan, Chesney Amy, Seal Shane, et al. Anatomical variations of the carpal tunnel structures. Canadian Journal of Plastic Surgery. 2009;17(3):3-7.
2. Börekci Ali, Selahi Özge, Tanriverdi Nursena, et al. Accessory hand muscles over the transverse carpal ligament: An obstacle in carpal tunnel surgery. World Neurosurgery. 2023;170:e402-e415.
3. Kapoor SK, Tiwari A, Kumar A, et al. Clinical relevance of palmaris longus agenesis: Common anatomical aberration. Anato Sci Int. 2008;83:45-48.
4. Frohse F, Fränkel M. Die muskeln des menschlichen armes, handbuch der anatomie des menschen. Munich & Jena, Gustav Fischer. 1908:115-118.
5. Reimann AF, Daseler EH, Anson BJ, Beaton LE. The palmaris longus muscle and tendon: A study of 1600 extremities. Anat Rec. 1944;89:495-505.
6. Bast Björn-Ole, Winkler Merle, Kurz Bodo. Reversed palmaris profundus muscle variation. Int J Anat Var (IJAV). 2016;9:21-24.
7. Pezas TP, Jose R. Palmaris profundus in the carpal tunnel. BMJ Case Reports CP. 2021;14:e241328.
8. Quinn T, White D. Palmaris profundus causing carpal tunnel syndrome: A rare anatomical variant. Aust J Plast Surg. 2019;2(2):71-73.

9. Sahinoglu Kayihan, Cassell Martin D, profundus). Annals of Anatomy-
Miyachi Ryosuke, et al. Musculus Anatomischer Anzeiger. 1994;176(3):229-
comitans nervi mediani (M. Palmaris 232.

© 2023 Baba et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97318>